



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Perlecan siRNA (m): sc-44394

BACKGROUND

Perlecan is part of a large family of heparan sulfate proteoglycans (HSPGs). As key components of cell surfaces and extracellular matrices, HSPGs modulate growth factor activities and thereby influence cell growth and differentiation. Additionally, HSPGs play a critical role in regulating tumor cell metastasis by mediating cell adhesion and the activities of growth and angiogenic factors. Perlecan consists of five distinct structural domains that interact with a number of matrix molecules, cytokines and growth factors to influence cartilage development and neuromuscular junction activity. Antithrombin, a key regulator of blood coagulation proteases, and TGF β 1 act as inhibitors and stimulators of Perlecan expression, respectively, interactions which may provide avenues for therapeutic intervention in certain types of cancer.

REFERENCES

- Hassell, J., Yamada, Y. and Arikawa-Hirasawa, E. 2002. Role of Perlecan in skeletal development and diseases. *Glycoconj. J.* 19: 263-267.
- Jiang, X. and Couchman, J.R. 2003. Perlecan and tumor angiogenesis. *J. Histochem. Cytochem.* 51: 1393-1410.
- Reiland, J., Sanderson, R.D., Waguespack, M., Barker, S.A., Long, R., Carson, D.D. and Marchetti, D. 2004. Heparanase degrades syndecan-1 and Perlecan Heparan sulfate: functional implications for tumor cell invasion. *J. Biol. Chem.* 279: 8047-8055.
- Zhang, W., Chuang, Y.J., Swanson, R., Li, J., Seo, K., Leung, L., Lau, L.F. and Olson, S.T. 2004. Antiangiogenic antithrombin downregulates the expression of the pro-angiogenic Heparan sulfate proteoglycan, Perlecan, in endothelial cells. *Blood* 103: 1185-1191.
- Casar, J.C., Cabello-Verrugio, C., Olguin, H., Aldunate, R., Inestrosa, N.C. and Brandan, E. 2004. Heparan sulfate proteoglycans are increased during skeletal muscle regeneration: requirement of Syndecan-3 for successful fiber formation. *J. Cell Sci.* 117: 73-84.

CHROMOSOMAL LOCATION

Genetic locus: Hspg2 (mouse) mapping to 4 D3.

PRODUCT

Perlecan siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Perlecan shRNA Plasmid (m): sc-44394-SH and Perlecan shRNA (m) Lentiviral Particles: sc-44394-V as alternate gene silencing products.

For independent verification of Perlecan (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44394A and sc-44394B.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Perlecan siRNA (m) is recommended for the inhibition of Perlecan expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Perlecan (A7L6): sc-33707 is recommended as a control antibody for monitoring of Perlecan gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Perlecan gene expression knockdown using RT-PCR Primer: Perlecan (m)-PR: sc-44394-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.