



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

eIF2C2 siRNA (h): sc-44409

BACKGROUND

Eukaryotic translation initiation factor 2C (eIF2C) proteins (argonaute family) influence RNA interference (RNAi) as components of the RNA-inducible silencing complex (RISC) or microRNA (miRNA)-containing ribonucleoprotein particle (miRNP). Small RNAs, including small interfering RNAs (siRNAs) and miRNAs, can silence target genes through mechanisms that utilize RISC or miRNP particles. eIF2C1 (argonaute 1, AGO1, eIF2C, GERP95, Q99) and Dicer1 play a coordinated role in siRNA-mediated gene silencing. eIF2C2 (slicer, argonaute 2, AGO2, Q10) is a RISC component that can concentrate in cytoplasmic processing bodies (P-bodies) and catalyze mRNA cleavage. Mammalian P-bodies contain mRNAs and have an association with miRNA-induced translational silencing and siRNA-induced mRNA degradation. Additional eIF2C proteins include eIF2C3 (argonaute 3, AGO3), eIF2C4 (argonaute 4, AGO4) and melf2c5 (mouse argonaute 5).

REFERENCES

- Martinez, J., et al. 2002. Single-stranded antisense siRNAs guide target RNA cleavage in RNAi. *Cell* 110: 563-574.
- Carmell, M.A., et al. 2002. The argonaute family: tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis. *Genes Dev.* 16: 2733-2742.
- Yan, K.S., et al. 2003. Structure and conserved RNA binding of the PAZ domain. *Nature* 426: 468-474.

CHROMOSOMAL LOCATION

Genetic locus: AGO2 (human) mapping to 8q24.3.

PRODUCT

eIF2C2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see eIF2C2 shRNA Plasmid (h): sc-44409-SH and eIF2C2 shRNA (h) Lentiviral Particles: sc-44409-V as alternate gene silencing products.

For independent verification of eIF2C2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44409A, sc-44409B and sc-44409C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

eIF2C2 siRNA (h) is recommended for the inhibition of eIF2C2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

eIF2C2 (4F9): sc-53521 is recommended as a control antibody for monitoring of eIF2C2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor eIF2C2 gene expression knockdown using RT-PCR Primer: eIF2C2 (h)-PR: sc-44409-PR (20 μ l, 431 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Aporntewan, C., et al. 2011. Hypomethylation of intragenic LINE-1 represses transcription in cancer cells through AGO2. *PLoS ONE* 6: e17934.
- Otsuka, M., et al. 2011. Receptor for activated protein kinase C: requirement for efficient microRNA function and reduced expression in hepatocellular carcinoma. *PLoS ONE* 6: e24359.
- Yoon, J.H., et al. 2012. LincRNA-p21 suppresses target mRNA translation. *Mol. Cell* 47: 648-655.
- Latorre, E., et al. 2016. Human antigen R binding and regulation of SOX2 mRNA in human mesenchymal stem cells. *Mol. Pharmacol.* 89: 243-252.
- Balaratnam, S., et al. 2018. A piRNA utilizes HILI and HIWI2 mediated pathway to down-regulate ferritin heavy chain 1 mRNA in human somatic cells. *Nucleic Acids Res.* 46: 10635-10648.
- Idda, M.L., et al. 2018. Cooperative translational control of polymorphic BAFF by NF90 and miR-15a. *Nucleic Acids Res.* 46: 12040-12051.
- Cross-Barnet, C., et al. 2019. Facilitators and barriers to optimal preventive service use among providers and older patients. *Geriatr. Nurs.* 40: 72-77.
- Tang, L., et al. 2019. Muscleblind-like 1 destabilizes Snail mRNA and suppresses the metastasis of colorectal cancer cells via the Snail/E-cadherin axis. *Int. J. Oncol.* 54: 955-965.
- Kulkarni, R.P., et al. 2019. miRNA-dependent regulation of STIM1 expression in breast cancer. *Sci. Rep.* 9: 13076.

RESEARCH USE

For research use only, not for use in diagnostic procedures.