



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Triadin shRNA (m) Lentiviral Particles: sc-44414-V

BACKGROUND

Triadin is a junctional terminal cisternae protein found mainly in human skeletal muscle. The gene TRDN which encodes for the protein maps to chromosome 6q22.31. Triadin, a type II membrane protein, is involved in anchoring calsequestrin to the sarcoplasmic reticulum, allowing its coupling with the ryanodine receptor (RyR). Triadin inhibits the calcium channel activity of ryanodine receptor in skeletal muscle. It co-localizes with RyR in the junctional sarcoplasmic reticulum membrane.

REFERENCES

- Caswell, A.H., et al. 1991. Localization and partial characterization of the oligomeric disulfide-linked molecular weight 95 kDa protein (Triadin) which binds the ryanodine and dihydropyridine receptors in skeletal muscle triadic vesicles. *Biochemistry* 30: 7507-7513.
- Flucher, B.E., et al. 1993. Triad formation: organization and function of the sarcoplasmic reticulum calcium release channel and Triadin in normal and dysgenic muscle *in vitro*. *J. Cell Biol.* 123: 1161-1174.
- Knudson, C.M., et al. 1993. Biochemical characterization of ultrastructural localization of a major junctional sarcoplasmic reticulum glycoprotein (Triadin). *J. Biol. Chem.* 268: 12637-12645.
- Knudson, C.M., et al. 1993. Primary structure and topological analysis of a skeletal muscle-specific junctional sarcoplasmic reticulum glycoprotein (Triadin). *J. Biol. Chem.* 268: 12646-12654.
- Brandt, N.R., et al. 1993. Detection and localization of triadin in rat ventricular muscle. *J. Membr. Biol.* 131: 219-228.
- Taske, N.L., et al. 1995. Molecular cloning of the cDNA encoding human skeletal muscle Triadin and its localisation to chromosome 6q22-6q23. *Eur. J. Biochem.* 233: 258-265.
- Ohkura, M., et al. 1998. Dual regulation of the skeletal muscle ryanodine receptor by Triadin and calsequestrin. *Biochemistry* 37: 12987-12993.

CHROMOSOMAL LOCATION

Genetic locus: Trdn (mouse) mapping to 10 A4.

PRODUCT

Triadin shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Triadin siRNA (m): sc-44414 and Triadin shRNA Plasmid (m): sc-44414-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80°C . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4°C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

Triadin shRNA (m) Lentiviral Particles is recommended for the inhibition of cardiac Triadin isoform 3 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Triadin (E-18): sc-33393 is recommended as a control antibody for monitoring of cardiac Triadin isoform 3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Triadin gene expression knockdown using RT-PCR Primer: Triadin (m)-PR: sc-44414-PR (20 μ l, 540 bp). Annealing temperature for the primers should be $55-60^\circ\text{C}$ and the extension temperature should be $68-72^\circ\text{C}$.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.