



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

INSIG-1 siRNA (m): sc-44433

BACKGROUND

INSIG-1 and INSIG-2 play distinct roles in a negative-feedback mechanism for cholesterol synthesis. INSIG-1 is highly expressed in liver and fibroblast cell lines. INSIG-1 localizes to the endoplasmic reticulum (ER) and binds the sterol-sensing domain of SREBP cleavage-activating protein (SCAP). Sterol induces INSIG-1 binding to SCAP. INSIG-2, another ER protein, binds SCAP in a sterol-regulated manner. Thus, INSIG-1 and INSIG-2 block the export of SCAP from the ER and ultimately inhibit cholesterol synthesis by preventing the proteolytic processing of SREBPs by Golgi enzymes. INSIG-1 is encoded by the Insulin-induced gene (INSIG-1). INSIG-1 gene expression is suppressed by oxysterols and restored following the introduction of the hypocholesterolemic agent LY295427. The negative feedback mechanism is absent in mutant CHO cells with a point mutation in one SCAP allele within the sterol-sensing domain. The mutant cells constitutively cleave SREBP in the presence of sterols. The critical role of INSIG-1 and INSIG-2 in cholesterol metabolism may be exploited as a therapeutic effect for hypercholesterolemia.

REFERENCES

- Peng, Y., et al. 1997. Cloning, human chromosomal assignment, and adipose and hepatic expression of the CL-6/INSIG1 gene. *Genomics* 43: 278-284.
- Janowski, B.A. 2002. The hypocholesterolemic agent LY295427 upregulates INSIG-1, identifying the INSIG-1 protein as a mediator of cholesterol homeostasis through SREBP. *Proc. Natl. Acad. Sci. USA* 99: 12675-12680.
- Yabe, D., et al. 2002. INSIG, a second endoplasmic reticulum protein that binds SCAP and blocks export of sterol regulatory element-binding proteins. *Proc. Natl. Acad. Sci. USA* 99: 12753-12758.
- Yang, T., et al. 2002. Crucial step in cholesterol homeostasis: sterols promote binding of SCAP to INSIG-1, a membrane protein that facilitates retention of SREBPs in ER. *Cell* 110: 489-500.
- Yabe, D., et al. 2002. Three mutations in sterol-sensing domain of SCAP block interaction with insig and render SREBP cleavage insensitive to sterols. *Proc. Natl. Acad. Sci. USA* 99: 16672-16677.

CHROMOSOMAL LOCATION

Genetic locus: *Insig1* (mouse) mapping to 5 B1.

PRODUCT

INSIG-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see INSIG-1 shRNA Plasmid (m): sc-44433-SH and INSIG-1 shRNA (m) Lentiviral Particles: sc-44433-V as alternate gene silencing products.

For independent verification of INSIG-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44433A, sc-44433B and sc-44433C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

INSIG-1 siRNA (m) is recommended for the inhibition of INSIG-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

INSIG-1 (A-9): sc-390504 is recommended as a control antibody for monitoring of INSIG-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor INSIG-1 gene expression knockdown using RT-PCR Primer: INSIG-1 (m)-PR: sc-44433-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.