



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# GSTM1 siRNA (h): sc-44461

## BACKGROUND

Members of the Glutathione S-transferase (GST) family of proteins function in the detoxification of xenobiotics to protect cells against toxicant-induced damage. GSTs are differentially expressed in lung, liver and kidney tissue and, notably, three isoforms (GSTA1-1, GSTA1-4 and GSTM1) localize to the mitochondria in addition to the cytoplasm. In normal and transformed cells, the oncoprotein Myb transcriptionally upregulates GSTM1. This isoform shows high specific activity for Aflatoxin B1 epoxide conjugation, suggesting an important role for this interaction in the defense against both chemical and oxidative stress.

## REFERENCES

- McGuire, S., et al. 1997. Increased levels of Glutathione S-transferases and appearance of novel  $\alpha$  class isoenzymes in kidneys of mice exposed to mercuric chloride. I. Biochemical and immunohistochemical studies. *Nephron* 77: 452-460.
- Massey, T.E., et al. 2000. Mechanisms of Aflatoxin B1 lung tumorigenesis. *Exp. Lung Res.* 26: 673-683.
- Raza, H., et al. 2002. Multiple isoforms of mitochondrial Glutathione S-transferases and their differential induction under oxidative stress. *Biochem. J.* 366: 45-55.
- Bartley, P.A., et al. 2003. Regulation of the gene encoding Glutathione S-transferase M1 (GSTM1) by the Myb oncoprotein. *Oncogene* 22: 7570-7575.
- McCarty, K.M., et al. 2006. Arsenic methylation, GSTT1, GSTM1, GSTP1 polymorphisms, and skin lesions. *Environ. Health Perspect.* 115: 341-345.
- Demir, A., et al. 2007. The role of GSTM1 gene polymorphisms in lung cancer development in Turkish population. *J. Carcinog.* 6: 13.
- Spurdle, A.B., et al. 2007. A systematic approach to analysing gene-gene interactions: polymorphisms at the microsomal epoxide hydrolase EPHX and Glutathione S-transferase GSTM1, GSTT1, and GSTP1 loci and breast cancer risk. *Cancer Epidemiol. Biomarkers Prev.* 16: 769-774.
- Moretti, M., et al. 2007. Primary DNA damage and genetic polymorphisms for CYP1A1, EPHX and GSTM1 in workers at a graphite electrode manufacturing plant. *BMC Public Health* 7: 270.

## CHROMOSOMAL LOCATION

Genetic locus: GSTM1 (human) mapping to 1p13.3.

## PRODUCT

GSTM1 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GSTM1 shRNA Plasmid (h): sc-44461-SH and GSTM1 shRNA (h) Lentiviral Particles: sc-44461-V as alternate gene silencing products.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

GSTM1 siRNA (h) is recommended for the inhibition of GSTM1 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

GSTM1 (1H4F2): sc-517262 is recommended as a control antibody for monitoring of GSTM1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GSTM1 gene expression knockdown using RT-PCR Primer: GSTM1 (h)-PR: sc-44461-PR (20  $\mu$ l, 556 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

- Kim, J.H., et al. 2012. Functional dissection of Nrf2-dependent phase II genes in vascular inflammation and endotoxic injury using Keap1 siRNA. *Free Radic. Biol. Med.* 53: 629-640.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.