



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



MGAT1 shRNA (h) Lentiviral Particles: sc-44467-V

BACKGROUND

Monoacylglycerol O-acyltransferase (MGAT) catalyzes diacylglycerol (a precursor to triacylglycerol) synthesis. MGAT is important in intestinal absorption of dietary fat because resynthesis of triacylglycerol is needed for the assembly of the lipoproteins that transport absorbed fat to tissues. MGAT1 is expressed in stomach, kidney, liver and adipose tissue but is not found in the intestine. On the contrary, MGAT2 (monoacylglycerol O-acyltransferase 2) is highly expressed in the small intestine as well as in kidney, liver, colon, stomach and white adipose tissue. MGAT 3 (monoacylglycerol O-acyltransferase 3) is highly homologous to MGAT1 and MGAT2. The expression of MGAT3 is restricted to the gastrointestinal tract, most concentrated in the ileum.

REFERENCES

1. Ioffe, E., et al. 1996. Essential role for complex N-glycans in forming an organized layer of bronchial epithelium. *Proc. Natl. Acad. Sci. USA* 93: 11041-11044.
2. Yip, B., et al. 1997. Organization of the human β -1,2-N-acetylglucosaminyltransferase I gene (MGAT1), which controls complex and hybrid N-glycan synthesis. *Biochem. J.* 321: 465-474.
3. Yen, C.L., et al. 2002. Identification of a gene encoding MGAT1, a monoacylglycerol acyltransferase. *Proc. Natl. Acad. Sci. USA* 99: 8512-8517.
4. Cheng, D., et al. 2003. Identification of acyl coenzyme A:monoacylglycerol acyltransferase 3, an intestinal specific enzyme implicated in dietary fat absorption. *J. Biol. Chem.* 278: 13611-13614.
5. Cao, J., et al. 2003. Cloning and functional characterization of a mouse intestinal acyl-CoA:monoacylglycerol acyltransferase, MGAT2. *J. Biol. Chem.* 278: 13860-13866.
6. Yen, C.L., et al. 2003. MGAT2, a monoacylglycerol acyltransferase expressed in the small intestine. *J. Biol. Chem.* 278: 18532-18537.
7. Shi, S., et al. 2004. Inactivation of the MGAT1 gene in oocytes impairs oogenesis, but embryos lacking complex and hybrid N-glycans develop and implant. *Mol. Cell. Biol.* 24: 9920-9929. Erratum in *Mol. Cell. Biol.* 25: 1214.

CHROMOSOMAL LOCATION

Genetic locus: MOGAT1 (human) mapping to 2q36.1.

PRODUCT

MGAT1 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see MGAT1 siRNA (h): sc-44467 and MGAT1 shRNA Plasmid (h): sc-44467-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80°C . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4°C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

MGAT1 shRNA (h) Lentiviral Particles is recommended for the inhibition of MGAT1 expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

MGAT1 (C-15): sc-32386 is recommended as a control antibody for monitoring of MGAT1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MGAT1 gene expression knockdown using RT-PCR Primer: MGAT1 (h)-PR: sc-44467-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ\text{C}$ and the extension temperature should be $68-72^\circ\text{C}$.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.