



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# NCX1 siRNA (m): sc-44515

## BACKGROUND

Sodium/calcium exchanger proteins are integral membrane proteins primarily seen in cardiac cells. In cardiac myocytes, the concentration of  $Ca^{2+}$  alternates between low levels during relaxation and high levels during contraction. The  $Na^+/Ca^{2+}$  exchanger 1 (NCX1) protein mediates  $Ca^{2+}$  extrusion from cardiac cells during relaxation. Four NCX1 isoforms (NCX1.1, NCX1.3, NCX1.7 and NCX1.10) result from alternate splicing. NCX1 mRNA is present at high levels in the heart, with lower levels present in the brain. NCX2 is most abundantly expressed in brain, in contrast the the broader distribution of NCX1, which is also expressed in heart, kidney, lung, smooth and skeletal muscle. The difference in expression for the transporter subtypes is believed to reflect differences in their functional roles. Regulation mechanisms for these exchanger proteins have not been fully characterized.

## REFERENCES

- Li, Z., et al. 1994. Cloning of the NCX2 isoform of the plasma membrane  $Na^+/Ca^{2+}$  exchanger. *J. Biol. Chem.* 269: 17434-17439.
- Kikuno, R., et al. 1999. Prediction of the coding sequences of unidentified human genes. XIV. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res.* 6: 197-205.
- Li, L., et al. 2000. Calcineurin controls the transcription of  $Na^+/Ca^{2+}$  exchanger isoforms in developing cerebellar neurons. *J. Biol. Chem.* 275: 20903-20910.
- Frayse, B., et al. 2001. Expression of the  $Na^+/Ca^{2+}$  exchanger in skeletal muscle. *Am. J. Physiol., Cell Physiol.* 280: C146-C154.
- Canitano, A., et al. 2002. Brain distribution of the  $Na^+/Ca^{2+}$  exchanger-encoding genes NCX1, NCX2, and NCX3 and their related proteins in the central nervous system. *Ann. N.Y. Acad. Sci.* 976: 394-404.
- Thurneysen, T., et al. 2002. Sodium/calcium exchanger subtypes NCX1, NCX2 and NCX3 show cell-specific expression in rat hippocampus cultures. *Brain. Res. Mol. Brain. Res.* 107: 145-156.
- Papa, M., et al. 2003. Differential expression of the  $Na^+/Ca^{2+}$  exchanger transcripts and proteins in rat brain regions. *J. Comp. Neurol.* 461: 31-48.
- Annunziato, L., et al. 2004. Pharmacology of brain  $Na^+/Ca^{2+}$  exchanger: from molecular biology to therapeutic perspectives. *Pharmacol. Rev.* 56: 633-654.
- SWISS-PROT/TrEMBL (Q9UPR5). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>.

## CHROMOSOMAL LOCATION

Genetic locus: Slc8a1 (mouse) mapping to 17 E3.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

NCX1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NCX1 shRNA Plasmid (m): sc-44515-SH and NCX1 shRNA (m) Lentiviral Particles: sc-44515-V as alternate gene silencing products.

For independent verification of NCX1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44515A, sc-44515B and sc-44515C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}$  C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}$  C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

NCX1 siRNA (m) is recommended for the inhibition of NCX1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NCX1 gene expression knockdown using RT-PCR Primer: NCX1 (m)-PR: sc-44515-PR (20  $\mu$ l, 449 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$  C and the extension temperature should be 68-72 $^{\circ}$  C.