



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# CD229 siRNA (h): sc-44551

## BACKGROUND

T lymphocyte surface antigen Ly9 (CD229), also designated lymphocyte antigen 9 or cell-surface molecule Ly9, is a cell surface glycoprotein. CD229 is a Type I membrane protein that is crucial in adhesion reactions between T lymphocytes and accessory cells (homophilic interaction). It belongs to the CD2 subfamily of the immunoglobulin gene superfamily of proteins (along with CD2, CD48, CD58, CD84, CD244 and CD150). Receptors of this family are important in cytokine production regulation and cytotoxicity of lymphocytes and NK cells. CD229 interacts with the SAP/SH2D1A protein. CD229 is expressed on mature B cells, T cells, thymocytes and NK cells.

## REFERENCES

- Sandrin, M.S., et al. 1992. Isolation and characterization of cDNA clones for mouse Ly9. *J. Immunol.* 149: 1636-1641.
- Sandrin, M.S., et al. 1996. Isolation and characterization of cDNA clones for Humly9: the human homologue of mouse Ly9. *Immunogenetics* 43: 13-19.
- Tovar, V., et al. 2000. Gene structure of the mouse leukocyte cell surface molecule Ly9. *Immunogenetics* 51: 788-793.
- Romero, X., et al. 2005. CD229 (Ly9) lymphocyte cell surface receptor interacts homophilically through its N-terminal domain and relocates to the immunological synapse. *J. Immunol.* 174: 7033-7042.
- Martin, M., et al. 2005. Identification of GRB2 as a novel binding partner of the signaling lymphocytic activation molecule-associated protein binding receptor CD229. *J. Immunol.* 174: 5977-5986.
- SWISS-PROT/TrEMBL (Q9HBG7). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

## CHROMOSOMAL LOCATION

Genetic locus: LY9 (human) mapping to 1q23.3.

## PRODUCT

CD229 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CD229 shRNA Plasmid (h): sc-44551-SH and CD229 shRNA (h) Lentiviral Particles: sc-44551-V as alternate gene silencing products.

For independent verification of CD229 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44551A, sc-44551B and sc-44551C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

CD229 siRNA (h) is recommended for the inhibition of CD229 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

CD229 (HLy 9.1.25): sc-58992 is recommended as a control antibody for monitoring of CD229 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CD229 gene expression knockdown using RT-PCR Primer: CD229 (h)-PR: sc-44551-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.