



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Myosin Ig siRNA (h): sc-44621

BACKGROUND

Actin is a highly conserved protein that is expressed in all eukaryotic cells. Actin filaments can form both stable and labile structures and are crucial components of microvilli and the contractile apparatus of muscle cells. Troponin facilitates interaction between Actin and Myosin by binding to Ca²⁺. Troponin is made up of at least two subunits, which are divergent in cardiac muscle, fast skeletal muscle and slow skeletal muscle. Myosin is a hexamer of two heavy chains (MHC) and four light chains (MLC) that interacts with Actin to generate the force for diverse cellular movements, including cytokinesis, phagocytosis and muscle contraction. MYO1G, the gene encoding for the Myosin Ig protein, maps to chromosome 7p13-

REFERENCES

1. Bárány, M. 1967. ATPase activity of Myosin correlated with speed of muscle shortening. *J. Gen. Physiol.* 50: 197-218.
2. Billeter, R., et al. 1980. Myosin types in human skeletal muscle fibers. *Histochemistry* 65: 249-259.
3. Whalen, R.G., et al. 1980. Contractile protein isozymes in muscle development: identification of an embryonic form of Myosin heavy chain. *Proc. Natl. Acad. Sci. USA* 76: 5197-5201.
4. Barton, P.J. and Buckingham, M.E. 1985. The Myosin alkali light chain proteins and their genes. *Biochem. J.* 231: 249-261.
5. Warrick, H.M. and Spudich, J.A. 1988. Myosin structure and function in cell motility. *Annu. Rev. Cell Biol.* 3: 379-421.
6. Hodge, T. and Cope, M.J. 2000. A Myosin family tree. *J. Cell Sci.* 113: 3353-3354.
7. Pierce, R.A., et al. 2001. The HA-2 minor histocompatibility antigen is derived from a diallelic gene encoding a novel human class I Myosin protein. *J. Immunol.* 167: 3223-3230.
8. Dumont, R.A., et al. 2002. Myosin I isozymes in neonatal rodent auditory and vestibular epithelia. *J. Assoc. Res. Otolaryngol.* 3: 375-389.

CHROMOSOMAL LOCATION

Genetic locus: MYO1G (human) mapping to 7p13.

PRODUCT

Myosin Ig siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Myosin Ig shRNA Plasmid (h): sc-44621-SH and Myosin Ig shRNA (h) Lentiviral Particles: sc-44621-V as alternate gene silencing products.

For independent verification of Myosin Ig (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44621A, sc-44621B and sc-44621C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Myosin Ig siRNA (h) is recommended for the inhibition of Myosin Ig expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Myosin Ig gene expression knockdown using RT-PCR Primer: Myosin Ig (h)-PR: sc-44621-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.