



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



# Dynein IC1, cytosolic shRNA (m) Lentiviral Particles: sc-44674-V

## BACKGROUND

Dyneins are multisubunit, high molecular weight ATPases that interact with microtubules to generate force by converting the chemical energy of ATP into the mechanical energy of movement. Cytoplasmic or axonemal Dynein heavy, intermediate, light and light-intermediate chains are all components of minus end-directed motors; the complex transports cellular cargos towards the central region of the cell. Axonemal Dynein motors contain one to three non-identical heavy chains and cause a sliding of microtubules in the axonemes of cilia and flagella in a mechanism necessary for cilia to beat and propel the cell. Cytoplasmic Dynein is an approximately 12 subunit complex of two heavy chains, two intermediate chains to anchor Dynein to its cargo, four smaller intermediate chains and several light chains. It performs functions necessary for cell survival such as organelle transport and centrosome assembly. The carboxy terminus of Dynein is important for microtubule-dependent motility and is highly conserved, while the amino terminal regions are more variable. Several proteins regulate Dynein activity, including dynactin, LIS1 and NudEL (NudE-like).

## REFERENCES

1. Mallik, R., et al. 2004. Cytoplasmic dynein functions as a gear in response to load. *Nature* 427: 649-652.
2. Malikov, V., et al. 2004. Cytoplasmic dynein nucleates microtubules to organize them into radial arrays *in vivo*. *Mol. Biol. Cell* 15: 2742-2749.
3. Asai, D.J., et al. 2004. The dynein heavy chain family. *J. Eukaryot. Microbiol.* 51: 23-29.
4. Li, J., et al. 2005. NudEL targets dynein to microtubule ends through LIS1. *Nat. Cell Biol.* 7: 686-690.
5. Seetharam, R.N., et al. 2005. High speed sliding of axonemal microtubules produced by outer arm dynein. *Cell Motil. Cytoskeleton* 60: 96-103.
6. Lee, W.L., et al. 2005. The offloading model for dynein function: differential function of motor subunits. *J. Cell Biol.* 168: 201-207.

## CHROMOSOMAL LOCATION

Genetic locus: *Dync1i1* (mouse) mapping to 6 A1.

## PRODUCT

Dynein IC1, cytosolic shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200  $\mu$ l frozen stock containing  $1.0 \times 10^6$  infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Dynein IC1, cytosolic siRNA (m): sc-44674 and Dynein IC1, cytosolic shRNA Plasmid (m): sc-44674-SH as alternate gene silencing products.

## STORAGE

Store lentiviral particles at  $-80^\circ$  C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at  $4^\circ$  C for up to one week. Avoid repeated freeze thaw cycles.

## APPLICATIONS

Dynein IC1, cytosolic shRNA (m) Lentiviral Particles is recommended for the inhibition of Dynein IC1, cytosolic expression in mouse cells.

## SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200  $\mu$ l frozen viral stock containing  $1.0 \times 10^6$  infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

## GENE EXPRESSION MONITORING

Dynein IC1, cytosolic (D-16): sc-32192 is recommended as a control antibody for monitoring of Dynein IC1, cytosolic gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Dynein IC1, cytosolic gene expression knockdown using RT-PCR Primer: Dynein IC1, cytosolic (m)-PR: sc-44674-PR (20  $\mu$ l). Annealing temperature for the primers should be  $55-60^\circ$  C and the extension temperature should be  $68-72^\circ$  C.

## BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

## RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.