



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ERR α siRNA (m): sc-44707

BACKGROUND

Estrogen related receptor α (ERR α) is a nuclear receptor in the superfamily of ligand-regulated transcription factors and is a member of the NR3B orphan nuclear receptor subgroup (consisting of α , β and γ). ERR α plays a role in modulating the estrogen signaling pathway. In addition, the expression of ERR α has been shown to increase during fasting and cold exposure. ERR α may be important for regulating mitochondrial biogenesis and oxidative metabolism by acting directly on genes necessary for mitochondrial function. Mice lacking ERR α are unable to maintain their body temperature in the cold. ERR α may also be involved in the maintenance and formation of cartilage. This information could be useful in finding therapeutic agents for a variety of diseases affecting the joints.

REFERENCES

- Chen, F., et al. 1999. Identification of two hERR2-related novel nuclear receptors utilizing bioinformatics and inverse PCR. *Gene* 228: 101-109.
- Hong, H., et al. 1999. Hormone-independent transcriptional activation and coactivator binding by novel orphan nuclear receptor ERR3. *J. Biol. Chem.* 274: 22618-22626.
- Greschik, H., et al. 2002. Structural and functional evidence for ligand-independent transcriptional activation by the estrogen-related receptor 3. *Mol. Cell* 9: 303-313.
- Hentschke, M., et al. 2003. Identification of PNRC2 and TLE1 as activation function-1 cofactors of the orphan nuclear receptor ERR γ . *Biochem. Biophys. Res. Commun.* 312: 975-982.

CHROMOSOMAL LOCATION

Genetic locus: Esrra (mouse) mapping to 19 A.

PRODUCT

ERR α siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ERR α shRNA Plasmid (m): sc-44707-SH and ERR α shRNA (m) Lentiviral Particles: sc-44707-V as alternate gene silencing products.

For independent verification of ERR α (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44707A, sc-44707B and sc-44707C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ERR α siRNA (m) is recommended for the inhibition of ERR α expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

ERR α (2ERR7): sc-65720 is recommended as a control antibody for monitoring of ERR α gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ERR α gene expression knockdown using RT-PCR Primer: ERR α (m)-PR: sc-44707-PR (20 μ l, 491 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

SELECT PRODUCT CITATIONS

- Zhang, X., et al. 2016. PGC-1 α /ERR α -Sirt3 pathway regulates DAergic neuronal death by directly deacetylating SOD2 and ATP synthase β . *Antioxid. Redox Signal.* 24: 312-328.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.