



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Glycophorin A shRNA (m) Lentiviral Particles: sc-44730-V

BACKGROUND

Glycophorins A, B and C are sialoglycoproteins of the human erythrocyte membrane, which bear the antigenic determinants for the MN, Ss and Gerbich blood groups, respectively. Glycophorins span the membrane once and present their amino-terminal end to the extracellular surface of the human erythrocyte. The genetic array of expressed glycophorin surface antigens on erythrocytes defines the blood group phenotype of the individual. The human Glycophorin A gene maps to chromosome 4q28.2-q31.1, contains 7 exons which are 97% homologous to Glycophorin B, and encodes a 150 amino acid protein. The human Glycophorin B gene maps to chromosome 4q28-q31 and encodes a 91 amino acid protein. The human Glycophorin C gene maps to chromosome 2q14-q21 and contains four exons. Glycophorin C transcript can generate two protein isoforms. Isoform 1 includes all 4 exons and encodes the full length 128 amino acid protein. Isoform 2 is missing exon 2 and encodes a 109 amino acid protein, which specifies the Yus subtype of the Gerbich phenotype.

REFERENCES

1. Andersson, L.C., et al. 1979. Glycophorin A as a cell surface marker of early erythroid differentiation in acute leukemia. *Int. J. Cancer* 23: 717-720.
2. Liszka, K., et al., 1983, Glycophorin A expression in malignant hematopoiesis. *Am. J. Hematol.* 15: 219-226.
3. Nakahata, T., et al. 1994. Cell surface antigen expression in human erythroid progenitors: erythroid and megakaryocytic markers. *Leuk. Lymphoma* 13: 401-409.
4. Sadahira, Y., et al. 1999. Immunohistochemical identification of erythroid precursors in paraffin embedded bone marrow sections: spectrin is a superior marker to glycophorin. *J. Clin. Pathol.* 52: 919-921.

CHROMOSOMAL LOCATION

Genetic locus: Gypa (mouse) mapping to 8 C2.

PRODUCT

Glycophorin A shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Glycophorin A siRNA (m): sc-44730 and Glycophorin A shRNA Plasmid (m): sc-44730-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

Glycophorin A shRNA (m) Lentiviral Particles is recommended for the inhibition of Glycophorin A expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Glycophorin A (I-20): sc-19453 is recommended as a control antibody for monitoring of Glycophorin A gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Glycophorin A gene expression knockdown using RT-PCR Primer: Glycophorin A (m)-PR: sc-44730-PR (20 μ l). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.