



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SEC14L2 shRNA (h) Lentiviral Particles: sc-44738-V

BACKGROUND

The monomeric, SEC14L2 (SEC14-like protein 2), also known as supernatant protein factor (SPF), α -tocopherol-associated protein or squalene transfer protein, functions as a carrier protein transferring tocopherols, as a transcriptional activator via its interaction with α -tocopherol and as a stimulator of conversion of microsomal squalene-2,3-oxide into lanosterol in cholesterol biosynthesis. High levels of SEC14L2 are expressed in liver, brain, intestine and prostate. Subcellular localization of SEC14L2 is cytoplasmic, but in the presence of α -tocopherol, SEC14L2 localizes in the nucleus. Activity of SEC14L2 depends on posttranslational modifications, specifically phosphorylation by PKA and PKC.

REFERENCES

1. Caras, I.W., et al. 1979. Effects of a supernatant protein activator on microsomal squalene-2,3-oxide-lanosterol cyclase. *J. Biol. Chem.* 254: 11816-11821.
2. Friedlander, E.J., et al. 1980. Supernatant protein factor facilitates intermembrane transfer of squalene. *J. Biol. Chem.* 255: 8042-8045.
3. Chin, J., et al. 1984. Role of supernatant protein factor and anionic phospholipid in squalene uptake and conversion by microsomes. *J. Biol. Chem.* 259: 11735-11738.
4. Shibata, N., et al. 2001. Supernatant protein factor, which stimulates the conversion of squalene to lanosterol, is a cytosolic squalene transfer protein and enhances cholesterol biosynthesis. *Proc. Natl. Acad. Sci. USA* 98: 2244-2249.
5. Singh, D.K., et al. 2003. Phosphorylation of supernatant protein factor enhances its ability to stimulate microsomal squalene monooxygenase. *J. Biol. Chem.* 278: 5646-5651.
6. Stocker, A., et al. 2003. Supernatant protein factor in complex with RRR- α -tocopherylquinone: a link between oxidized Vitamin E and cholesterol biosynthesis. *J. Mol. Biol.* 332: 759-765.

CHROMOSOMAL LOCATION

Genetic locus: SEC14L2 (human) mapping to 22q12.2.

PRODUCT

SEC14L2 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see SEC14L2 siRNA (h): sc-44738 and SEC14L2 shRNA Plasmid (h): sc-44738-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80°C . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4°C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

SEC14L2 shRNA (h) Lentiviral Particles is recommended for the inhibition of SEC14L2 expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

SEC14L2 (H-4): sc-271905 is recommended as a control antibody for monitoring of SEC14L2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgM-HRP: sc-2064 (dilution range: 1:500-1:5,000), TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgM-FITC: sc-2082 (dilution range: 1:100-1:400) or goat anti-mouse IgM-TR: sc-2983 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SEC14L2 gene expression knockdown using RT-PCR Primer: SEC14L2 (h)-PR: sc-44738-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ\text{C}$ and the extension temperature should be $68-72^\circ\text{C}$.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.