



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Apelin shRNA (m) Lentiviral Particles: sc-44742-V

BACKGROUND

Apelin (APEL), an endogenous ligand for APJ, is an alternate co-receptor with CD4 for HIV-1 infection. This secreted protein inhibits HIV-1 entry into cells that coexpress APJ and CD4. By proteolytic processing of the peptide precursor, several different active peptides may be produced. Apelin-36, one such inotropic peptide, is being investigated as a potential plasma marker of cardiopulmonary disease. Apelin is highly expressed in brain, mainly in the thalamus, frontal cortex, hypothalamus and midbrain. Apelin is also secreted by the mammary gland into the colostrum and milk. Oral intake of Apelin (in milk and colostrum) may be important in the modulation of the immune responses in neonates and newborns. Apelin has also been found to be a potent stimulator of cardiac contractility and may function in the regulation of the cardiovascular system.

REFERENCES

1. Tatemoto, K., et al. 1998. Isolation and characterization of a novel endogenous peptide ligand for the human APJ receptor. *Biochem. Biophys. Res. Commun.* 251: 471-476.
2. Habata, Y., et al. 1999. Apelin, the natural ligand of the orphan receptor APJ, is abundantly secreted in the colostrum. *Biochim. Biophys. Acta* 1452: 25-35.
3. Lee, D.K., et al. 2000. Characterization of Apelin, the ligand for the APJ receptor. *J. Neurochem.* 74: 34-41.
4. Cayabyab, M., et al. 2000. Apelin, the natural ligand of the orphan seven-transmembrane receptor APJ, inhibits human immunodeficiency virus type 1 entry. *J. Virol.* 74: 11972-11976.
5. Wei, L., et al. 2005. Regulation of Apelin mRNA expression by Insulin and glucocorticoids in mouse 3T3-L1 adipocytes. *Regul. Pept.* 132: 27-32.

CHROMOSOMAL LOCATION

Genetic locus: *Apln* (mouse) mapping to X A4.

PRODUCT

Apelin shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Apelin siRNA (m): sc-44742 and Apelin shRNA Plasmid (m): sc-44742-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

Apelin shRNA (m) Lentiviral Particles is recommended for the inhibition of Apelin expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Apelin (S-20): sc-33469 is recommended as a control antibody for monitoring of Apelin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Apelin gene expression knockdown using RT-PCR Primer: Apelin (m)-PR: sc-44742-PR (20 μ l, 540 bp). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.