



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

BST-1 siRNA (m): sc-44751

BACKGROUND

BST-1 (bone marrow stromal antigen-1) has been identified as a surface molecule that is GPI-anchored to the cell membrane of stromal cells. Both ADP-ribosyl cyclase and cADPR hydrolase activities have been demonstrated by BST-1. cADPR activity is a potential regulator of Insulin secretion in pancreatic β cells. Most pancreatic islet cells express BST-1, indicating a link between BST-1 and Insulin secretion. BST-1 expression has also been found in a wide range of tissues including umbilical vein endothelial cells, monocytes and granulocytes. BST-1 expression in thymus tissue and on B and T cell progenitors undergoing gene rearrangement implicates BST-1 as a useful marker for lymphoid progenitor cells initiation gene rearrangement of their antigen receptors. BST-1 has also been shown to facilitate B cell growth and may act as a receptor.

REFERENCES

1. Kaisho, T., Ishikawa, J., Oritani, K., Inazawa, J., Tomizawa, H., Muraoka, O., Ochi, T. and Hirano, T. 1994. BST-1, a surface molecule of bone marrow stromal cell lines that facilitates pre-B-cell growth. *Proc. Natl. Acad. Sci. USA* 91: 5325-5329.
2. Hirata, Y., Kimura, N., Sato, K., Ohsugi, Y., Takasawa, S., Okamoto, H., Ishikawa, J., Kaisho, T., Ishihara, K. and Hirano, T. 1994. ADP ribosyl cyclase activity of a novel bone marrow stromal cell surface molecule, BST-1. *FEBS Lett.* 356: 244-248.
3. Kato, I., Takasawa, S., Akabane, A., Tanaka, O., Abe, H., Takamura, T., Suzuki, Y., Nata, K., Yonekura, H. and Yoshimoto, T. 1995. Regulatory role of CD38 (ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase) in Insulin secretion by glucose in pancreatic β cells. Enhanced Insulin secretion in CD38-expressing transgenic mice. *J. Biol. Chem.* 270: 30045-30050.
4. Kajimoti, Y., Miyagawa, J., Ishihara, K., Okuyama, Y., Fujitani, Y., Itoh, M., Yoshida, H., Kaisho, T., Matsuoka, T., Watada, H., Hanafusa, T., Yamasaki, Y., Kamada, T., Matsuzawa, Y. and Hirano, T. 1996. Pancreatic islet cells express BST-1, a CD38-like surface molecule having ADP-ribosyl cyclase activity. *Biochem. Biophys. Res. Commun.* 219: 941-946.
5. Okuyama, Y., Ishihara, K., Kimura, N., Hirata, Y., Sato, K., Itoh, M., Ok, L.B. and Hirano, T. 1996. Human BST-1 expressed on myeloid cells functions as a receptor molecule. *Biochem. Biophys. Res. Commun.* 228: 838-845.

CHROMOSOMAL LOCATION

Genetic locus: Bst1 (mouse) mapping to 5 B3.

PRODUCT

BST-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see BST-1 shRNA Plasmid (m): sc-44751-SH and BST-1 shRNA (m) Lentiviral Particles: sc-44751-V as alternate gene silencing products.

For independent verification of BST-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44751A, sc-44751B and sc-44751C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

BST-1 siRNA (m) is recommended for the inhibition of BST-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor BST-1 gene expression knockdown using RT-PCR Primer: BST-1 (m)-PR: sc-44751-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.