



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



FBP2 shRNA (h) Lentiviral Particles: sc-44831-V

BACKGROUND

Activation of FUSE, the far-upstream element, is required for the proper expression of the mammalian gene *c-Myc* in undifferentiated cells. The binding of FBP (FUSE-binding protein or Far upstream element binding protein) to FUSE is necessary for *c-Myc* expression, indicating that FBP functions as a growth-dependent regulator of *c-Myc* expression. Isolated from proliferating HL60 cells, FBP, FBP2, and FBP3 comprise a family of single-stranded DNA-binding proteins that specifically bind to FUSE elements. The FBP transcription factors share a conserved central DNA-binding domain and show significant homology in their carboxyl-terminal activation domains. Expression of FBP is detected in undifferentiated cells and is substantially decreased following cellular differentiation.

REFERENCES

- Avigan, M.I., et al. 1990. A far upstream element stimulates *c-Myc* expression in undifferentiated leukemia cells. *J. Biol. Chem.* 265: 18538-18545.
- Duncan, R.D., et al. 1994. A sequence-specific, single strand binding protein activates the far upstream of *c-Myc* and defines a new DNA binding motif. *Genes Dev.* 8: 465-480.
- Bazar, L., et al. 1995. A transactivator of *c-Myc* is coordinately regulated with the proto-oncogene during cellular growth. *Oncogene* 10: 2229-2238.
- Davis-Smyth, T., et al. 1996. The far upstream element-binding proteins comprise an ancient family of single-strand DNA-binding transactivators. *J. Biol. Chem.* 271: 31679-31687.
- Michelotti, G.A., et al. 1996. Multiple single-stranded cis elements are associated with activated chromatin of the human *c-Myc* gene *in vivo*. *Mol. Cell. Biol.* 16: 2656-2669.
- Rehbein, M., et al. 2002. Molecular characterization of MARTA1, a protein interacting with the dendritic targeting element of MAP2 mRNAs. *J. Neurochem.* 82: 1039-1046

CHROMOSOMAL LOCATION

Genetic locus: KHSRP (human) mapping to 19p13.3.

PRODUCT

FBP2 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see FBP2 siRNA (h): sc-44831 and FBP2 shRNA Plasmid (h): sc-44831-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80°C . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4°C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

FBP2 shRNA (h) Lentiviral Particles is recommended for the inhibition of FBP2 expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

FBP2 (D-12): sc-33031 is recommended as a control antibody for monitoring of FBP2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FBP2 gene expression knockdown using RT-PCR Primer: FBP2 (h)-PR: sc-44831-PR (20 μ l, 564 bp). Annealing temperature for the primers should be $55-60^\circ\text{C}$ and the extension temperature should be $68-72^\circ\text{C}$.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.