



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CAT-1 siRNA (h): sc-44923

BACKGROUND

The cationic amino acid transporter (CAT) family of proteins are part of a larger superfamily, the amino acid-polyamine-organocation (APC) superfamily. High-affinity cationic amino acid transporter-1 (CAT-1), also designated ecotropic retroviral leukemia receptor homolog, ATRC1 or REC1L, is an ubiquitously expressed integral membrane protein. In non-hepatic tissues, CAT-1 acts as a high-affinity, low capacity permease that is important in cationic amino acid transport. CAT-1 is also a potential ecotropic retroviral leukemia receptor. SLC7A1, the gene encoding for the CAT-1 protein, maps to chromosome 13q12.3.

REFERENCES

1. Yoshimoto, T., et al. 1991. Molecular cloning and characterization of a novel human gene homologous to the murine ecotropic retroviral receptor. *Virology* 185: 10-17.
2. Albritton, L.M., et al. 1992. The human cationic amino acid transporter (ATRC1): physical and genetic mapping to 13q12-q14. *Genomics* 12: 430-434.
3. Kamath, S.G., et al. 1999. Identification of three cationic amino acid transporters in placental trophoblast: cloning, expression and characterization of hCAT-1. *J. Membr. Biol.* 171: 55-62.
4. Zani, B.G. and Bohlen, H.G. 2005. Transport of extracellular L-arginine via cationic amino acid transporter is required during *in vivo* endothelial nitric oxide production. *Am. J. Physiol. Heart Circ. Physiol.* 289: H1381-H1390.

CHROMOSOMAL LOCATION

Genetic locus: SLC7A1 (human) mapping to 13q12.3.

PRODUCT

CAT-1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CAT-1 shRNA Plasmid (h): sc-44923-SH and CAT-1 shRNA (h) Lentiviral Particles: sc-44923-V as alternate gene silencing products.

For independent verification of CAT-1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44923A, sc-44923B and sc-44923C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CAT-1 siRNA (h) is recommended for the inhibition of CAT-1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

CAT-1 (F-2): sc-515782 is recommended as a control antibody for monitoring of CAT-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CAT-1 gene expression knockdown using RT-PCR Primer: CAT-1 (h)-PR: sc-44923-PR (20 μ l, 446 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Mandal, A., et al. 2017. L-arginine uptake by cationic amino acid transporter promotes intra-macrophage survival of *Leishmania donovani* by enhancing arginase-mediated polyamine synthesis. *Front. Immunol.* 8: 839.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.