

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Bcl-11a shRNA (m) Lentiviral Particles: sc-44935-V



The Power to Overtion

BACKGROUND

Bcl-11a (CtIP-1, EVI9, B cell CLL/lymphoma 11A) and Bcl-11b (CtIP-2, Rit1, B cell CLL/lymphoma 11B) genes play crucial roles in lymphopoiesis and influence the progression of hematopoietic malignancies. Disruption of the Bcl-11b locus is linked to T cell acute lymphoblastic leukemia and the loss of heterozygosity in mice results in T cell lymphoma. Bcl-11 proteins are related $\rm C_2H_2$ zinc-finger transcription factors that act as transcriptional repressors. Bcl-11b can interact with the metastasis-associated proteins MTA1 and MTA2 through the amino-terminal region. Bcl-11a is essential for postnatal development and normal lymphopoiesis. The Bcl-11a mouse gene is a common site of retroviral integration in myeloid leukemia, and may function as a leukemia disease gene, in part, through its interaction with Bcl-6.

REFERENCES

- 1. Dyer, M.J., et al. 2002. The configuration of the immunoglobulin genes in B cell chronic lymphocytic leukemia. Leukemia 16: 973-984.
- Avram, D., et al. 2002. COUP-TF (chicken ovalbumin upstream promoter transcription factor)-interacting protein-1 (CtIP-1) is a sequence-specific DNA binding protein. Biochem. J. 368: 555-563.
- Durum, S.K. 2003. Bcl-11: sibling rivalry in lymphoid development. Nat. Immunol. 4: 512-514.
- Nagel, S., et al. 2003. The cardiac homeobox gene Nkx2.5 is deregulated by juxtaposition with Bcl-11b in pediatric T-ALL cell lines via a novel t(5;14)(q35.1;q32.2). Cancer Res. 63: 5329-5334.
- Togashi, T., et al. 2004. Two distinct methods analyzing chromatin structure using centrifugation and antibodies to modified Histone H3: both provide similar chromatin states of the Rit1/Bcl-11b gene. Biochem. Biophys. Res. Commun. 313: 489-495.
- Sakata, J., et al. 2004. Involvement of V(D)J recombinase in the generation of intragenic deletions in the Rit1/Bcl-11b tumor suppressor gene in γ-ray-induced thymic lymphomas and in normal thymus of the mouse. Carcinogenesis 25: 1069-1075.

CHROMOSOMAL LOCATION

Genetic locus: Bcl11a (mouse) mapping to 11 A3.2.

PRODUCT

Bcl-11a shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μl frozen stock containing 1.0 x 10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Bcl-11a siRNA (m): sc-44935 and Bcl-11a shRNA Plasmid (m): sc-44935-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

Bcl-11a shRNA (m) Lentiviral Particles is recommended for the inhibition of Bcl-11a expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0 x 10 6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Bcl-11a (H-116): sc-366825 is recommended as a control antibody for monitoring of Bcl-11a gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Bcl-11a gene expression knockdown using RT-PCR Primer: Bcl-11a (m)-PR: sc-44935-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**