



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Bcl-11a siRNA (m): sc-44935



The Power to Question

BACKGROUND

Bcl-11a (CtIP-1, EVI9, B cell CLL/lymphoma 11A) and Bcl-11b (CtIP-2, Rit1, B cell CLL/lymphoma 11B) genes play crucial roles in lymphopoiesis and influence the progression of hematopoietic malignancies. Disruption of the Bcl-11b locus is linked to T cell acute lymphoblastic leukemia and the loss of heterozygosity in mice results in T cell lymphoma. Bcl-11 proteins are related C₂H₂ zinc-finger transcription factors that act as transcriptional repressors. Bcl-11b can interact with the metastasis-associated proteins MTA1 and MTA2 through the amino-terminal region. Bcl-11a is essential for postnatal development and normal lymphopoiesis. The Bcl-11a mouse gene is a common site of retroviral integration in myeloid leukemia, and may function as a leukemia disease gene, in part, through its interaction with Bcl-6.

REFERENCES

- Dyer, M.J., et al. 2002. The configuration of the immunoglobulin genes in B cell chronic lymphocytic leukemia. *Leukemia* 16: 973-984.
- Avram, D., et al. 2002. COUP-TF (chicken ovalbumin upstream promoter transcription factor)-interacting protein-1 (CtIP-1) is a sequence-specific DNA binding protein. *Biochem. J.* 368: 555-563.
- Durum, S.K. 2003. Bcl-11: sibling rivalry in lymphoid development. *Nat. Immunol.* 4: 512-514.
- Liu, P., et al. 2003. Bcl-11a is essential for normal lymphoid development. *Nat. Immunol.* 4: 525-532.
- Nagel, S., et al. 2003. The cardiac homeobox gene Nkx2.5 is deregulated by juxtaposition with Bcl-11b in pediatric T-ALL cell lines via a novel t(5;14)(q35.1;q32.2). *Cancer Res.* 63: 5329-5334.
- Togashi, T., et al. 2004. Two distinct methods analyzing chromatin structure using centrifugation and antibodies to modified Histone H3: both provide similar chromatin states of the Rit1/Bcl-11b gene. *Biochem. Biophys. Res. Commun.* 313: 489-495.

CHROMOSOMAL LOCATION

Genetic locus: Bcl11a (mouse) mapping to 11 A3.2.

PRODUCT

Bcl-11a siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Bcl-11a shRNA Plasmid (m): sc-44935-SH and Bcl-11a shRNA (m) Lentiviral Particles: sc-44935-V as alternate gene silencing products.

For independent verification of Bcl-11a (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44935A, sc-44935B and sc-44935C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Bcl-11a siRNA (m) is recommended for the inhibition of Bcl-11a expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Bcl-11a (H-1): sc-514842 is recommended as a control antibody for monitoring of Bcl-11a gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG_k BP-HRP: sc-516102 or m-IgG_k BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG_k BP-FITC: sc-516140 or m-IgG_k BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Bcl-11a gene expression knockdown using RT-PCR Primer: Bcl-11a (m)-PR: sc-44935-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.