



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



# BAI-3 shRNA (h) Lentiviral Particles: sc-45212-V

## BACKGROUND

Brain-specific angiogenesis inhibitors, including BAI-1, BAI-2 and BAI-3, are integral membrane proteins belonging to the G protein-coupled receptor 2 family. In addition to inhibiting angiogenesis in the brain, BAI proteins are also expressed in the heart, thymus, skeletal muscle and a variety of cell lines. Expression patterns of BAI-2 in the developing brain distinctly increase as development progresses. Localization of BAI-2 in most neurons of the cerebral cortex is analogous with BAI-1 expression. The activity of brain-specific angiogenesis inhibitor proteins has been inversely correlated with vascularization in some cancer tissues. Ischemic brain shows notably decreased expression of BAI-2 after hypoxia, which precludes increased expression of vascular endothelial growth factor, VEGF. A better understanding of the anti-angiogenic activity of these BAIs may offer potential therapeutic benefits.

## REFERENCES

1. Nishimori, H., et al. 1997. A novel brain-specific p53-target gene, BAI-1, containing thrombospondin type 1 repeats, inhibits experimental angiogenesis. *Oncogene* 15: 2145-2150.
2. Shiratsuchi, T., et al. 1998. Cloning and characterization of BAI-associated protein 1: a PDZ domain-containing protein that interacts with BAI-1. *Biochemistry* 247: 597-604.
3. Kee, H.J., et al. 2002. Expression of brain-specific angiogenesis inhibitor 2 (BAI-2) in normal and ischemic brain: involvement of BAI-2 in the ischemia-induced brain. *J. Cereb. Blood Flow Metab.* 22: 1054-1067.
4. Kaur, B., et al. 2003. Brain angiogenesis inhibitor 1 is differentially expressed in normal brain and glioblastoma independently of p53 expression. *Am. J. Pathol.* 162: 19-27.

## CHROMOSOMAL LOCATION

Genetic locus: BAI3 (human) mapping to 6q12.

## PRODUCT

BAI-3 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200  $\mu$ l frozen stock containing  $1.0 \times 10^6$  infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see BAI-3 siRNA (h): sc-45212 and BAI-3 shRNA Plasmid (h): sc-45212-SH as alternate gene silencing products.

## STORAGE

Store lentiviral particles at  $-80^\circ\text{C}$ . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at  $4^\circ\text{C}$  for up to one week. Avoid repeated freeze thaw cycles.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

BAI-3 shRNA (h) Lentiviral Particles is recommended for the inhibition of BAI-3 expression in human cells.

## SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200  $\mu$ l frozen viral stock containing  $1.0 \times 10^6$  infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

## GENE EXPRESSION MONITORING

BAI-3 (V-19): sc-33467 is recommended as a control antibody for monitoring of BAI-3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor BAI-3 gene expression knockdown using RT-PCR Primer: BAI-3 (h)-PR: sc-45212-PR (20  $\mu$ l). Annealing temperature for the primers should be  $55-60^\circ\text{C}$  and the extension temperature should be  $68-72^\circ\text{C}$ .

## BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

## RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.