



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## TEF-4 siRNA (m): sc-45233

### BACKGROUND

The transcriptional enhancer factor (TEF)/TEAD family includes TEF-1, TEF-3, TEF-4 and TEF-5. These proteins share a highly conserved 68 amino acid TEA/ATTS DNA-binding domain which binds to SV40 GT-IIC (GGAATG), SphI (AGTATG), SphII (AGCATG) and muscle-specific M-CAT (GGTATG) enhancers. TEFs are differentially expressed in human cultured cell lines and mouse embryonic and extra-embryonic tissues. Specifically, TEF-4 is strongly co-expressed with TEF-1 in mouse mitotic neuroblasts, and is also detected in the gut and the nephrogenic region of the kidney. TEF-4 associates with the powerful transcriptional coactivator Yap65 to mediate mitogenic signals. In addition, TEF-4 promotes the activation of the CTP:phosphocholine cytidyltransferase (CCT)  $\alpha$  protein, which is the rate-limiting enzyme of phosphatidylcholine biosynthesis by enhancing the transcriptional activity of Ets-1.

### REFERENCES

- Jacquemin, P., et al. 1996. A novel family of developmentally regulated mammalian transcription factors containing the TEA/ATTS DNA-binding domain. *J. Biol. Chem.* 271: 21775-21785.
- Jacquemin, P., et al. 1999. Localization of human transcription factor TEF-4 and TEF-5 (TEAD2, TEAD3) genes to chromosomes 19q13.3 and 6p21.2 using fluorescence *in situ* hybridization and radiation hybrid analysis. *Genomics* 55: 127-129.
- Jiang, S.W., et al. 2000. Cooperative binding of TEF-1 to repeated GGAATG-related consensus elements with restricted spatial separation and orientation. *DNA Cell. Biol.* 19: 507-514.
- Sugimoto, H., et al. 2001. Identification of transcriptional enhancer factor-4 as a transcriptional modulator of CTP:phosphocholine cytidyltransferase  $\alpha$ . *J. Biol. Chem.* 276: 12338-12344.
- Vassilev, A., et al. 2001. TEAD/TEF transcription factors utilize the activation domain of Yap65, a Src/Yes-associated protein localized in the cytoplasm. *Genes Dev.* 15: 1229-1241.

### CHROMOSOMAL LOCATION

Genetic locus: Tead2 (mouse) mapping to 7 B4.

### PRODUCT

TEF-4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TEF-4 shRNA Plasmid (m): sc-45233-SH and TEF-4 shRNA (m) Lentiviral Particles: sc-45233-V as alternate gene silencing products.

For independent verification of TEF-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45233A, sc-45233B and sc-45233C.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

TEF-4 siRNA (m) is recommended for the inhibition of TEF-4 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TEF-4 gene expression knockdown using RT-PCR Primer: TEF-4 (m)-PR: sc-45233-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.