



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

IL-18R siRNA (h): sc-45285

BACKGROUND

Interleukin-18 (IL-18) has been identified as a molecule that induces IFN- γ production and enhances NK cell cytotoxicity. IL-18 receptor (IL-18R) is a type I membrane protein present in lung, leukocytes, spleen, liver, thymus, prostate, small intestine, colon, placenta and heart, while absent from brain, skeletal muscle, pancreas and kidney. IL-18R is present in Hodgkin's disease cell lines, and does not bind IL-1 α or IL-1 β . The association of IL-18 to IL-18R leads to activation of NF κ B. At present two subunits of IL-18R have been characterized: IL-18R α and IL-18R β . IL-18R α has been described as the ligand-binding chain and IL-18R β as the signal-transduction chain.

REFERENCES

1. Torigoe, K., Ushio, S., Okura, T., Kobayashi, S., Taniai, M., Kunikata, T., Murakami, T., Sanou, O., Kojima, H., Fujii, M., Ohta, T., Ikeda, M., Ikegami, H. and Kurimoto, M. 1997. Purification and characterization of the human interleukin-18 receptor. *J. Biol. Chem.* 272: 25737-25742.
2. Strand, M.L., Wahlgren, A., Svechnikov, K., Zetterström, C., Setchell, B.P. and Söder, O. 2005. Interleukin-18 is expressed in rat testis and may promote germ cell growth. *Mol. Cell. Endocrinol.* 240: 64-73.
3. Vermot-Desroches, C., Subiger, O., Guenot, F., Sergent, E., Bonnin, B. and Wijdenes, J. 2005. Monoclonal antibodies specific for the IL-18 receptor. *Cell. Immunol.* 236: 101-104.
4. Zhou, Y., Yamaguchi, E., Fukui, Y., Konno, S., Maeda, Y., Kimata, K. and Nishimura, M. 2005. Enhanced expression of interleukin-18 receptor α chain by CD4⁺ T cells in sarcoidosis. *Chest* 128: 2497-2503.
5. Krasna, E., Kolesár, L., Slavcev, A., Valhová, S., Kronosová, B., Jaresová, M. and Stráz, I. 2005. IL-18 receptor expression on epithelial cells is upregulated by TNF α . *Inflammation* 29: 33-37.
6. Asakawa, M., Kono, H., Amemiya, H., Matsuda, M., Suzuki, T., Maki, A. and Fujii, H. 2006. Role of interleukin-18 and its receptor in hepatocellular carcinoma associated with hepatitis C virus infection. *Int. J. Cancer* 118: 564-570.

CHROMOSOMAL LOCATION

Genetic locus: IL18R1 (human) mapping to 2q12.1.

PRODUCT

IL-18R siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IL-18R shRNA Plasmid (h): sc-45285-SH and IL-18R shRNA (h) Lentiviral Particles: sc-45285-V as alternate gene silencing products.

For independent verification of IL-18R (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45285A, sc-45285B and sc-45285C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IL-18R siRNA (h) is recommended for the inhibition of IL-18R expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IL-18R gene expression knockdown using RT-PCR Primer: IL-18R (h)-PR: sc-45285-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Mo, C., Dai, Y., Kang, N., Cui, L. and He, W. 2012. Ectopic expression of human MutS homologue 2 on renal carcinoma cells is induced by oxidative stress with interleukin-18 promotion via p38 mitogen-activated protein kinase (MAPK) and c-Jun N-terminal kinase (JNK) signaling pathways. *J. Biol. Chem.* 287: 19242-19254.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.