



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



synphilin-1 shRNA (m) Lentiviral Particles: sc-45293-V

BACKGROUND

Synphilin-1 (α -synuclein interacting protein, SNCAIP) is a 919-amino acid protein that associates with α -synuclein and promotes the formation of cytosolic inclusions in neuronal cells. The synuclein family members, including α -synuclein and β -synuclein, are predominantly expressed in the brain where they influence synaptic regulation and neuronal plasticity. Synphilin-1 contains modular protein domains, such as ankyrin-like repeats and a coiled-coil domain. While both α -synuclein and synphilin-1 are co-expressed in Lewy bodies of patients with Parkinson's disease (PD), only mutations in the gene for α -synuclein have been determined to confer pathogenicity.

REFERENCES

1. Ueda, K., et al. 1993. Molecular cloning of cDNA encoding an unrecognized component of amyloid in Alzheimer disease. *Proc. Natl. Acad. Sci. USA* 90: 11282-11286.
2. Jakes, R., et al. 1994. Identification of two distinct synucleins from human brain. *FEBS Letts.* 345: 27-32.
3. Engelender, S., et al. 1999. Synphilin-1 associates with α -synuclein and promotes the formation of cytosolic inclusions. *Nat. Genet.* 22: 110-114.
4. Wakabayashi, K., et al. 2000. Synphilin-1 is present in Lewy bodies in Parkinson's disease. *Ann. Neurol.* 47: 521-553.
5. Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 603779. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Kawamata, H., et al. 2001. Interaction of α -synuclein and synphilin-1: effect of Parkinson's disease-associated mutations. *J. Neurochem.* 77: 929-934.

CHROMOSOMAL LOCATION

Genetic locus: Sncap (mouse) mapping to 18 D1.

PRODUCT

synphilin-1 shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see synphilin-1 siRNA (m): sc-45293 and synphilin-1 shRNA Plasmid (m): sc-45293-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

synphilin-1 shRNA (m) Lentiviral Particles is recommended for the inhibition of synphilin-1 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

synphilin-1 (H-300): sc-98296 is recommended as a control antibody for monitoring of synphilin-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor synphilin-1 gene expression knockdown using RT-PCR Primer: synphilin-1 (m)-PR: sc-45293-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ$ C and the extension temperature should be $68-72^\circ$ C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.