



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AMPK α 1/2 siRNA (h): sc-45312

BACKGROUND

AMPK (for 5'-AMP-activated protein kinase) is a heterotrimeric complex comprising a catalytic α subunit and regulatory β and γ subunits. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. AMPK is activated by high AMP and low ATP through a mechanism involving allosteric regulation, promotion of phosphorylation by an upstream protein kinase known as AMPK kinase, and inhibition of dephosphorylation. Activated AMPK can phosphorylate and regulate *in vivo* hydroxymethylglutaryl-CoA reductase and acetyl-CoA carboxylase, which are key regulatory enzymes of sterol synthesis and fatty acid synthesis, respectively. The human AMPK α 1 and AMPK α 2 genes encode 548 amino acid and 552 amino acid proteins, respectively. Human AMPK β 1 encodes a 271 amino acid protein and human AMPK β 2 encodes a 272 amino acid protein. The human AMPK γ 1 gene encodes a 331 amino acid protein. Human AMPK γ 2 and AMPK γ 3, which are 569 and 492 amino acid proteins, respectively, contain unique N-terminal domains and may participate directly in the binding of AMP within the AMPK complex.

CHROMOSOMAL LOCATION

Genetic locus: PRKAA1 (human) mapping to 5p13.1, PRKAA2 (human) mapping to 1p32.2.

PRODUCT

AMPK α 1/2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AMPK α 1/2 shRNA Plasmid (h): sc-45312-SH and AMPK α 1/2 shRNA (h) Lentiviral Particles: sc-45312-V as alternate gene silencing products.

For independent verification of AMPK α 1/2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45312A, sc-45312B and sc-45312C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AMPK α 1/2 siRNA (h) is recommended for the inhibition of AMPK α 1/2 expression in human cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

AMPK α 1/2 (D-6): sc-74461 is recommended as a control antibody for monitoring of AMPK α 1/2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

SELECT PRODUCT CITATIONS

- Cao, C., et al. 2008. AMP-activated protein kinase contributes to UV- and H₂O₂-induced apoptosis in human skin keratinocytes. *J. Biol. Chem.* 283: 28897-28908.
- Lin, C.L., et al. 2008. Epigallocatechin gallate (EGCG) attenuates high glucose-induced Insulin signaling blockade in human Hep G2 hepatoma cells. *Mol. Nutr. Food Res.* 52: 930-939.
- Schneider, A., et al. 2008. Hypoxia-induced energy stress inhibits the mTOR pathway by activating an AMPK/REDD1 signaling axis in head and neck squamous cell carcinoma. *Neoplasia* 10: 1295-1302.
- Park, J.S., et al. 2019. Repositioning of niclosamide ethanolamine (NEN), an anthelmintic drug, for the treatment of lipotoxicity. *Free Radic. Biol. Med.* 137: 143-157.
- Wei, Q., et al. 2019. Maslinic acid inhibits colon tumorigenesis by the AMPK-mTOR signaling pathway. *J. Agric. Food Chem.* 67: 4259-4272.
- He, L., et al. 2019. PINK1/Parkin-mediated mitophagy promotes apelin-13-induced vascular smooth muscle cell proliferation by AMPK α and exacerbates atherosclerotic lesions. *J. Cell. Physiol.* 234: 8668-8682.
- Jung, T.W., et al. 2019. Protectin DX ameliorates palmitate-induced hepatic insulin resistance through AMPK/SIRT1-mediated modulation of fetuin-A and SeP expression. *Clin. Exp. Pharmacol. Physiol.* 46: 898-909.
- Kazyken, D., et al. 2019. AMPK directly activates mTORC2 to promote cell survival during acute energetic stress. *Sci. Signal.* 12 pii: eaav3249.
- Lee, J., et al. 2019. Metformin induces apoptosis and inhibits proliferation through the AMP-activated protein kinase and Insulin-like growth factor 1 receptor pathways in the bile duct cancer cells. *J. Cancer* 10: 1734-1744.

RESEARCH USE

For research use only, not for use in diagnostic procedures.