



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

# IRAK-4 siRNA (m): sc-45401

## BACKGROUND

Interleukin-1 receptor (IL1R)-associated kinases (IRAKs) are important mediators in the signal transduction of Toll-like receptor (TLR) and IL1R family members, collectively referred to as TIRs. IRAK family members include two active kinases, IRAK-1 and IRAK-4, and two inactive kinases, IRAK-2 and IRAK-M. Binding of IL-1 to its cognate receptor results in the activation of the NF $\kappa$ B signaling pathway and MAP kinase pathways. IRAK-4 appears to act up-stream of other IRAKs and phosphorylates IRAK-1 on threonine 387. It is highly expressed in liver and kidney tissues, but also displays a wide, low level of expression in other tissues. IRAK-4 is an essential component of innate immunity. Deficiency of IRAK-4 leads to recurrent bacterial infections and profound hyporesponsiveness to LPS and IL-1. Therefore, IRAK-4 may be a potential target for therapeutic drug design.

## REFERENCES

- Li, S., et al. 2002. IRAK-4: a novel member of the IRAK family with the properties of an IRAK-kinase. *Proc. Natl. Acad. Sci. USA* 99: 5567-5572.
- Janssens, S., et al. 2003. Functional diversity and regulation of different interleukin-1 receptor-associated kinase (IRAK) family members. *Mol. Cell* 11: 293-302.
- Lye, E., et al. 2004. The role of interleukin 1 receptor-associated kinase-4 (IRAK-4) kinase activity in IRAK-4-mediated signaling. *J. Biol. Chem.* 279: 40653-40658.
- Medvedev, A.E., et al. 2005. Cutting edge: expression of IL-1 receptor-associated kinase-4 (IRAK-4) proteins with mutations identified in a patient with recurrent bacterial infections alters normal IRAK-4 interaction with components of the IL-1 receptor complex. *J. Immunol.* 174: 6587-6591.
- Lasker, M.V., et al. 2005. Cutting edge: molecular structure of the IL-1R-associated kinase-4 death domain and its implications for TLR signaling. *J. Immunol.* 175: 4175-4179.

## CHROMOSOMAL LOCATION

Genetic locus: *Irak4* (mouse) mapping to 15 E3.

## PRODUCT

IRAK-4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IRAK-4 shRNA Plasmid (m): sc-45401-SH and IRAK-4 shRNA (m) Lentiviral Particles: sc-45401-V as alternate gene silencing products.

For independent verification of IRAK-4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45401A, sc-45401B and sc-45401C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

IRAK-4 siRNA (m) is recommended for the inhibition of IRAK-4 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IRAK-4 gene expression knockdown using RT-PCR Primer: IRAK-4 (m)-PR: sc-45401-PR (20  $\mu$ l, 491 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

- Wang, Y.Y., et al. 2016. Increased translocation of antigens to endosomes and TLR4 mediated endosomal recruitment of TAP contribute to nicotine augmented cross-presentation. *Oncotarget* 7: 38451-38466.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.