



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

plexin-B2 siRNA (m): sc-45423

BACKGROUND

Plexins are a family of large, transmembrane receptors for multiple classes of semaphorins in vertebrates. They are widely expressed, and regions of their extracellular domain are homologous to both scatter factor receptors and semaphorin domains. Plexins may act as semaphorin receptors alone or in combination with neuropilins. Plexins are divided into four subfamilies designated plexin-A, -B, -C, and -D. Plexin-B1 and -B2 are both receptors for SEMA4D, which stimulates axonal outgrowth of embryonic dorsal root ganglion neurons. Plexin-B3 binds to SEMA5A, which controls axon guidance and can initiate the intracellular signaling of the hepatocyte growth factor/scatter factor receptor Met.

REFERENCES

1. Artigiani, S., et al. 1992. Plexins, semaphorins, and scatter factor receptors: a common root for cell guidance signals? *IUBMB Life* 48: 477-478.
2. Kolodkin, A.L., et al. 1993. The semaphorin genes encode a family of transmembrane and secreted growth cone guidance molecules. *Cell* 75: 1389-1399.
3. Kameyama, T., et al. 1996. Identification of plexin family molecules in mice. *Biochem. Biophys. Res. Commun.* 226: 396-402.
4. Tamagnone, L., et al. 1997. Control of invasive growth by hepatocyte growth factor (HGF) and related scatter factors. *Cytokine Growth Factor Rev.* 8: 129-142.
5. Winberg, M.L., et al. 1998. Plexin-A is a neuronal semaphorin receptor that controls axon guidance. *Cell* 95: 903-916.
6. Takahashi, T., et al. 1999. Plexin-neuropilin-1 complexes form functional semaphorin-3A receptors. *Cell* 99: 59-69.
7. Tamagnone, L., et al. 1999. Plexins are a large family of receptors for transmembrane, secreted and GPI-anchored semaphorins in vertebrates. *Cell* 99: 71-80.
8. Masuda, K. et al. 2004. Sema4D stimulates axonal outgrowth of embryonic DRG sensory neurones. *Genes Cells* 9: 821-829.
9. Artigiani, S. et al. 2004. Plexin-B3 is a functional receptor for semaphorin 5A. *EMBO Rep.* 5: 710-714.

CHROMOSOMAL LOCATION

Genetic locus: Plxn2 (mouse) mapping to 15 E3.

PRODUCT

plexin-B2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see plexin-B2 shRNA Plasmid (m): sc-45423-SH and plexin-B2 shRNA (m) Lentiviral Particles: sc-45423-V as alternate gene silencing products.

For independent verification of plexin-B2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45423A, sc-45423B and sc-45423C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

plexin-B2 siRNA (m) is recommended for the inhibition of plexin-B2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

plexin-B2 (A-5): sc-373969 is recommended as a control antibody for monitoring of plexin-B2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor plexin-B2 gene expression knockdown using RT-PCR Primer: plexin-B2 (m)-PR: sc-45423-PR (20 μ l, 533 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.