



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ARP siRNA (h): sc-45435

BACKGROUND

The gene encoding arginine-rich protein (ARP), also designated ARMET, which is highly conserved in all species, localizes to human chromosome 3p21.2. Mutation of ARP occurs in several human tumors, including primary head and neck, non-small-cell lung, renal cell, breast and prostate cancers. Previously, malignancy of the ARP gene was thought to be the result of frequent variations of the triplet AGG repeat around codon 50, but studies showed no significant difference in this variation between normal and cancer patient populations. Subsequently, it has been shown that the ARP protein contains a smaller N-terminal region, which does not include the arginine-rich region, and that codon 50 actually is the start codon for the protein. A function for the ARP protein has yet to be determined.

REFERENCES

- Shridhar, V., Rivard, S., Shridhar, R., Mullins, C., Bostick, L., Sakr, W., Grignon, D., Miller, O.J. and Smith, D.I. 1996. A gene from human chromosomal band 3p21.1 encodes a highly conserved arginine-rich protein and is mutated in renal cell carcinomas. *Oncogene* 12: 1931-1939.
- Evron, E., Cairns, P., Halachmi, N., Ahrendt, S.A., Reed, A.L. and Sidransky, D. 1997. Normal polymorphism in the incomplete trinucleotide repeat of the arginine-rich protein gene. *Cancer Res.* 57: 2888-2889.
- Shridhar, V., Rivard, S., Wang, X., Shridhar, R., Paisley, C., Mullins, C., Beinrat, L., Dugan, M., Sarkar, F., Miller, O.J., Vaitkevicius, V.K. and Smith, D.I. 1997. Mutations in the arginine-rich protein gene (ARP) in pancreatic cancer. *Oncogene* 14: 2213-2216.
- Tanaka, H., Shimada, Y., Harada, H., Shinoda, M., Hatooka, S., Imamura, M. and Ishizaki, K. 2000. Polymorphic variation of the ARP gene on 3p21 in Japanese esophageal cancer patients. *Oncol. Rep.* 7: 591-593.
- Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 601916. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: MANF (human) mapping to 3p21.2.

PRODUCT

ARP siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ARP shRNA Plasmid (h): sc-45435-SH and ARP shRNA (h) Lentiviral Particles: sc-45435-V as alternate gene silencing products.

For independent verification of ARP (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45435A, sc-45435B and sc-45435C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ARP siRNA (h) is recommended for the inhibition of ARP expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

ARP (B-3): sc-515906 is recommended as a control antibody for monitoring of ARP gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ARP gene expression knockdown using RT-PCR Primer: ARP (h)-PR: sc-45435-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.