



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Matrilin-2 siRNA (h): sc-45447

BACKGROUND

The matrilin family proteins are secreted extracellular matrix proteins. Matrilin-1 is a homotrimer that binds to collagen and is a component of the extracellular matrix of nonarticular cartilage. It is secreted primarily by chondrocytes in a characteristic spatial, temporal and developmental stage-specific pattern during skeletogenesis. Matrilin-2 is a secreted protein involved in matrix assembly. Matrilin-3 is a secreted protein expressed solely in cartilaginous tissues. It is important in the extracellular matrix of cartilage and in the formation of extracellular filamentous networks. Matrilin-4, expressed in embryonic kidney, lung and placenta, is a secreted protein important to the extracellular matrix of cartilage.

REFERENCES

1. Deak, F., et al. 1999. The matrilins: a novel family of oligomeric extracellular matrix proteins. *Matrix Biol.* 18: 55-64.
2. Strusberg, I., et al. 2002. Association analysis of genotypic frequencies of Matrilin-1 gene in patients with osteoarthritis. *Clin. Exp. Rheumatol.* 20: 543-545.
3. Ohno, S., et al. 2003. Immunohistochemical study of Matrilin-1 in arthritic articular cartilage of the mandibular condyle. *J. Oral Pathol. Med.* 32: 237-242.
4. Wiberg, C., et al. 2003. Complexes of Matrilin-1 and Biglycan or Decorin connect Collagen VI microfibrils to both Collagen II and aggrecan. *J. Biol. Chem.* 278: 37698-37704.
5. Mann, H.H., et al. 2004. Interactions between the cartilage oligomeric matrix protein and matrilins. Implications for matrix assembly and the pathogenesis of chondrodysplasias. *J. Biol. Chem.* 279: 25294-25298.

CHROMOSOMAL LOCATION

Genetic locus: MATN2 (human) mapping to 8q22.1.

PRODUCT

Matrilin-2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Matrilin-2 shRNA Plasmid (h): sc-45447-SH and Matrilin-2 shRNA (h) Lentiviral Particles: sc-45447-V as alternate gene silencing products.

For independent verification of Matrilin-2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45447A, sc-45447B and sc-45447C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Matrilin-2 siRNA (h) is recommended for the inhibition of Matrilin-2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Matrilin-2 (2B8): sc-293397 is recommended as a control antibody for monitoring of Matrilin-2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Matrilin-2 gene expression knockdown using RT-PCR Primer: Matrilin-2 (h)-PR: sc-45447-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.