

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# TIEG1 shRNA (h) Lentiviral Particles: sc-45463-V



The Power to Overtion

#### **BACKGROUND**

Originally isolated from osteoblastic cells, the TGF $\beta$ -inducible early gene-1 (TIEG1) is a Kruppel-like zinc-finger transcription factor-encoding gene which regulates cellular growth and differentiation. TIEG1 is regulated as an early response gene by TGF $\beta$ 1. It is expressed in both acinar and ductular epithelial cells from exocrine pancreas and may serve as an early response gene in pancreatic cell lines. Further, overexpression of TIEG1 in TGF $\beta$ -sensitive epithelial cells induces apoptosis. TIEG1 and EGR $\alpha$  are expressed from alternate promoters of the same gene. Both are highly expressed in human fetal osteoblast cells. TIEG1 is additionally expressed at high levels in PBLs, spleen and colon, and at lower levels in thymus, small intestine, ovary, prostate and skeletal muscle. The nuclear TIEG2 protein, which shares significant homology with TIEG1, was originally isolated from globin-expressing human fetal erythroid cells. TIEG2 is also expressed in fetal liver. Overexpression of TIEG2 in cultured epithelial cells inhibits cellular proliferation. TIEG2 expression is upregulated by TGF $\beta$ 1 and serum deprivation.

#### **REFERENCES**

- Subramaniam, M., et al. 1995. Identification of a novel TGFβ-regulated gene encoding a putative zinc-finger protein in human osteoblasts. Nucleic Acids Res. 23: 4907-4912.
- Blok, L.J., et al. 1995. Characteriza-tion of an early growth response gene, which encodes a zinc-finger transcription factor, potentially involved in cell cycle regulation. Mol. Endocrinol. 9: 1610-1620.
- Tachibana, I., et al. 1997. Overexpression of the TGFβ-regulated zinc-finger encoding gene, TIEG, induces apoptosis in pancreatic epithelial cells. J. Clin. Invest. 99: 2365-2374.

#### CHROMOSOMAL LOCATION

Genetic locus: KLF10 (human) mapping to 8q22.3.

#### **PRODUCT**

TIEG1 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 µl frozen stock containing 1.0 x 10<sup>6</sup> infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see TIEG1 siRNA (h): sc-45463 and TIEG1 shRNA Plasmid (h): sc-45463-SH as alternate gene silencing products.

#### **STORAGE**

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

#### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

#### **APPLICATIONS**

TIEG1 shRNA (h) Lentiviral Particles is recommended for the inhibition of TIEG1 expression in human cells.

#### **SUPPORT REAGENTS**

Control shRNA Lentiviral Particles: sc-108080. Available as 200  $\mu$ l frozen viral stock containing 1.0 x 10 $^6$  infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

#### **GENE EXPRESSION MONITORING**

TIEG1 (H-190): sc-67062 is recommended as a control antibody for monitoring of TIEG1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **RT-PCR REAGENTS**

Semi-quantitative RT-PCR may be performed to monitor TIEG1 gene expression knockdown using RT-PCR Primer: TIEG1 (h)-PR: sc-45463-PR (20  $\mu$ I, 558 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

#### **BIOSAFETY**

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

#### **RESEARCH USE**

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**