



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TIEG1 shRNA (m) Lentiviral Particles: sc-45464-V

BACKGROUND

Originally isolated from osteoblastic cells, the TGF β -inducible early gene-1 (TIEG1) is a Krüppel-like zinc finger transcription factor-encoding gene which regulates cellular growth and differentiation. TIEG1 is regulated as an early response gene by TGF β 1. It is expressed in both acinar and ductular epithelial cells from exocrine pancreas and may serve as an early response gene in pancreatic cell lines. Further, overexpression of TIEG1 in TGF β -sensitive epithelial cells induces apoptosis. TIEG1 and EGR α are expressed from alternate promoters of the same gene. Both are highly expressed in human fetal osteoblast cells. TIEG1 is additionally expressed at high levels in PBLs, spleen and colon, and at lower levels in thymus, small intestine, ovary, prostate and skeletal muscle. The nuclear TIEG2 protein, which shares significant homology with TIEG1, was originally isolated from globin-expressing human fetal erythroid cells. TIEG2 is also expressed in fetal liver. Overexpression of TIEG2 in cultured epithelial cells inhibits cellular proliferation. TIEG2 expression is upregulated by TGF β 1 and serum deprivation.

REFERENCES

- Subramaniam, M., et al. 1995. Identification of a novel TGF- β -regulated gene encoding a putative zinc finger protein in human osteoblasts. *Nucleic Acids Res.* 23: 4907-4912.
- Blok, L.J., et al. 1995. Characterization of an early growth response gene, which encodes a zinc finger transcription factor, potentially involved in cell cycle regulation. *Mol. Endocrinol.* 9: 1610-1620.
- Tachibana, I., et al. 1997. Overexpression of the TGF β -regulated zinc finger encoding gene, TIEG, induces apoptosis in pancreatic epithelial cells. *J. Clin. Invest.* 99: 2365-2374.

CHROMOSOMAL LOCATION

Genetic locus: Klf10 (mouse) mapping to 15 B3.1.

PRODUCT

TIEG1 shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see TIEG1 siRNA (m): sc-45464 and TIEG1 shRNA Plasmid (m): sc-45464-SH as alternate gene silencing products.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

APPLICATIONS

TIEG1 shRNA (m) Lentiviral Particles is recommended for the inhibition of TIEG1 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

TIEG1 (C-14): sc-34544 is recommended as a control antibody for monitoring of TIEG1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TIEG1 gene expression knockdown using RT-PCR Primer: TIEG1 (m)-PR: sc-45464-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.