



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

RANTES siRNA (m): sc-45573

BACKGROUND

Structurally, C-C or β -chemokines are characterized by a set of conserved, adjacent cysteines. Members of this family include MCP-1, MCP-2, MCP-3, MIP-1 α , MIP-1 β , RANTES and I-309. RANTES (regulated upon activation, normal T cell expressed and secreted) is expressed by platelets, eosinophils, fibroblasts, macrophages, endothelial cells and T lymphocytes. Consistent with its belonging to the chemokine family, RANTES exhibits strong chemoattractant activity towards monocytes and NK cells. I-309 was initially identified as a factor present in γ/δ T lymphocytes. I-309 cDNA encodes a protein 73 amino acids in length with one potential N-linked glycosylation site. Unlike the other members of the C-C family, I-309 does not induce chemotaxis in natural killer (NK) cells.

REFERENCES

1. Miller, M.D., et al. 1989. A novel polypeptide secreted by activated human T lymphocytes. *J. Immunol.* 143: 2907-2916.
2. Lloyd, A.R., et al. 1996. Chemokines regulate T cell adherence to recombinant adhesion molecules and extracellular matrix proteins. *J. Immunol.* 156: 932-938.
3. Wells, T.N., et al. 1996. Selectivity and antagonism of chemokine receptors. *J. Leukoc. Biol.* 59: 53-60.
4. Taub, D.D., et al. 1996. β chemokines costimulate lymphocyte cytolysis, proliferation and lymphokine production. *J. Leukoc. Biol.* 59: 81-89.
5. Wang, J.H., et al. 1996. Expression of RANTES by human bronchial epithelial cells *in vitro* and *in vivo* and the effect of corticosteroids. *Am. J. Respir. Cell Mol. Biol.* 14: 27-35.
6. Ying, S., et al. 1996. Human eosinophils express messenger RNA encoding RANTES and store and release biologically active RANTES protein. *Eur. J. Immunol.* 26: 70-76.
7. Loetscher, P., et al. 1996. Activation of NK cells by C-C chemokines. Chemotaxis, Ca^{2+} mobilization and enzyme release. *J. Immunol.* 156: 322-327.

CHROMOSOMAL LOCATION

Genetic locus: Ccl5 (mouse) mapping to 11 C.

PRODUCT

RANTES siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RANTES shRNA Plasmid (m): sc-45573-SH and RANTES shRNA (m) Lentiviral Particles: sc-45573-V as alternate gene silencing products.

For independent verification of RANTES (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45573A and sc-45573B.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RANTES siRNA (m) is recommended for the inhibition of RANTES expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

RANTES (A-4): sc-365826 is recommended as a control antibody for monitoring of RANTES gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RANTES gene expression knockdown using RT-PCR Primer: RANTES (m)-PR: sc-45573-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.