



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Selenoprotein R siRNA (m): sc-45575

BACKGROUND

Selenium is an essential trace element that is incorporated as selenocysteine into the primary structure of selenoproteins. Nutritional deficiency of selenium decreases selenoprotein concentrations and leads to pathologic conditions. Most of the known selenoproteins are members of the glutathione peroxidase or iodothyronine deiodinase families. Selenoprotein P (SEPP1) is a major selenoprotein that is not a member of those families. It is an extracellular glycoprotein that is present in several isoforms and is the only selenoprotein known to contain multiple selenocysteine residues. SECIS elements form stem-loop structures in the 3' untranslated regions (UTR) of eukaryotic mRNAs that encode selenoproteins. The Selenoprotein W SECIS elements contain an additional highly conserved base-paired stem that may prevent inappropriate selenocysteine incorporation at the UGA stop codons. Selenoprotein R, also designated methionine-R-sulfoxide reductase or Selenoprotein X1, is a zinc-binding protein. The gene encoding for Selenoprotein R contains selenocysteine, which is encoded by the usual stop codon TGA, at the active site. Selenoprotein R may play a role in protection against oxidative stress.

REFERENCES

- Hill, K.E., et al. 1993. Conserved nucleotide sequences in the open reading frame and 3' untranslated region of Selenoprotein P mRNA. *Proc. Natl. Acad. Sci. USA* 90: 537-541.
- Chittum, H.S., et al. 1996. Multiple forms of Selenoprotein P in rat plasma. *Arch. Biochem. Biophys.* 325: 124-128.
- Hill, K.E., et al. 1996. Human Selenoprotein P gene maps to 5q31. *Genomics* 36: 550-551.
- Kryukov, G.V., et al. 1999. New mammalian selenocysteine-containing proteins identified with an algorithm that searches for selenocysteine insertion sequence elements. *J. Biol. Chem.* 274: 33888-33897.
- Kryukov, G.V., et al. 2002. Selenoprotein R is a zinc-containing stereo-specific methionine sulfoxide reductase. *Proc. Natl. Acad. Sci. USA* 99: 4245-4250.

CHROMOSOMAL LOCATION

Genetic locus: *Sepr1* (mouse) mapping to 17 A3.3.

PRODUCT

Selenoprotein R siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Selenoprotein R shRNA Plasmid (m): sc-45575-SH and Selenoprotein R shRNA (m) Lentiviral Particles: sc-45575-V as alternate gene silencing products.

For independent verification of Selenoprotein R (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45575A, sc-45575B and sc-45575C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Selenoprotein R siRNA (m) is recommended for the inhibition of Selenoprotein R expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Selenoprotein R gene expression knockdown using RT-PCR Primer: Selenoprotein R (m)-PR: sc-45575-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.