



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Acrosin shRNA (m) Lentiviral Particles: sc-45605-V

BACKGROUND

Acrosin, a member of the peptidase S1 family, is a major protease present in the acrosome of mature mammalian spermatozoa. Acrosin is a typical serine proteinase with trypsin-like cleavage specificity. The zymogen form, proacrosin, is the precursor of Acrosin synthesized only in the postmeiotic stages of spermatogenesis. The active enzyme functions in the lysis of the zona pellucida, allowing the penetration of sperm through the innermost glycoprotein layers of the ovum.

REFERENCES

1. Peknicova, J., et al. 1990. Monoclonal antibodies against boar acrosomal antigens labelling undamaged acrosomes of spermatozoa in immunofluorescence test. *Andrologia* 22: 427-435.
2. Moos, J. et al. 1993. Protein-protein interactions controlling acrosin release and solubilization during the boar sperm acrosome reaction. *Biol. Reprod.* 49: 408-415.
3. Peknicova, J., et al. 2001. Monoclonal antibodies to intra-acrosomal proteins inhibit gamete binding *in vitro*. *Theriogenology* 56: 211-223.
4. Wang, H., et al. 2004. Novel role for a sterol response element binding protein in directing spermatogenic cell-specific gene expression. *Mol. Cell. Biol.* 24: 10681-10688.
5. Chaudhury, K., et al. 2004. Studies on the membrane integrity of human sperm treated with a new injectable male contraceptive. *Hum. Reprod.* 19: 1826-1830.
6. Pyoung, I., et al. 2004. Molecular characterization of the 32 kDa boar sperm protease. *Mol. Reprod. Dev.* 68: 354-358.
7. Zalata, A.A., et al. 2004. Relationship between acrosin activity of human spermatozoa and oxidative stress. *Asian J. Androl.* 6: 313-318.
8. Langlois, M.R., et al. 2005. Discrepancy between sperm acrosin activity and sperm morphology: significance for fertilization *in vitro*. *Clin. Chim. Acta* 351: 121-129.

CHROMOSOMAL LOCATION

Genetic locus: Acr (mouse) mapping to 15 E3.

PRODUCT

Acrosin shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Acrosin siRNA (m): sc-45605 and Acrosin shRNA Plasmid (m): sc-45605-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

Acrosin shRNA (m) Lentiviral Particles is recommended for the inhibition of Acrosin expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Acrosin (C-14): sc-46284 is recommended as a control antibody for monitoring of Acrosin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Acrosin gene expression knockdown using RT-PCR Primer: Acrosin (m)-PR: sc-45605-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ$ C and the extension temperature should be $68-72^\circ$ C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.