



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FcRH1 shRNA (h) Lentiviral Particles: sc-45628-V

BACKGROUND

The Fc receptor homolog (FcRH) family of proteins are related to the classical Fc receptors (FcR) and belong to the immunoglobulin receptor superfamily. The proteins in the FcRH family are type I transmembrane glycoproteins and consist of FcRH1-FcRH6. The gene encoding for the proteins maps to chromosome 1q23.1, near its FcR relatives. The FcRH proteins, which are involved in immune system regulation, have immunoreceptor-tyrosine inhibitory motifs in their cytoplasmic domains. Mutations in the gene encoding for the FcRH proteins may be associated with systemic lupus erythematosus, auto-immune thyroid disease and rheumatoid arthritis. The FcRH genes are expressed primarily, although not exclusively, by mature B lineage cells, and may serve important regulatory roles in normal and neoplastic B cell development.

REFERENCES

1. Davis, R.S., et al. 2001. Identification of a family of Fc receptor homologs with preferential B cell expression. *Proc. Natl. Acad. Sci. USA* 98: 9772-9777.
2. Davis, R.S., et al. 2002. Fc receptor homologs (FcRH1-5) extend the Fc receptor family. *Curr. Top. Microbiol. Immunol.* 266: 85-112
3. Davis, R.S., et al. 2002. Fc receptor homologs: newest members of a remarkably diverse Fc receptor gene family. *Immunol. Rev.* 190: 123-136.
4. Davis, R.S., et al. 2004. Differential B cell expression of mouse Fc receptor homologs. *Int. Immunol.* 16: 1343-1353.
5. Davis, R.S., et al. 2005. An extended family of Fc receptor relatives. *Eur. J. Immunol.* 35: 674-680.
6. Leu, C.M., et al. 2005. FcRH1: an activation coreceptor on human B cells. *Blood* 105: 1121-1126.

CHROMOSOMAL LOCATION

Genetic locus: FCRL1 (human) mapping to 1q23.1.

PRODUCT

FcRH1 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see FcRH1 siRNA (h): sc-45628 and FcRH1 shRNA Plasmid (h): sc-45628-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

FcRH1 shRNA (h) Lentiviral Particles is recommended for the inhibition of FcRH1 expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

FcRH1 (E-8): sc-515003 is recommended as a control antibody for monitoring of FcRH1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FcRH1 gene expression knockdown using RT-PCR Primer: FcRH1 (h)-PR: sc-45628-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ$ C and the extension temperature should be $68-72^\circ$ C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.