



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FcRn siRNA (m): sc-45633

BACKGROUND

FcRn present in the intestinal epithelium of neonatal mice and rats mediates the selective uptake of immunoglobulin G (IgG) in mothers' milk, thereby helping newborn animals to acquire passive immunity. FcRn (also designated FCGRT, Brambell receptor, FcRn α chain, IgG Gc receptor and neonatal Fc receptor) is comprised of a heavy chain and β -2-Microglobulin. FcRn heavy chain shows approximately 35% amino acid identity to an MHC class I molecule. FcRn localizes in endosomes of vascular endothelial cells and selectively recycles IgG to the cell surface, thus protecting IgG from lysosomal catabolism. This protection mechanism is a major constituent for ensuring IgG are the longest lived of all plasma proteins.

REFERENCES

1. Claypool, S.M., et al. 2002. Functional reconstitution of human FcRn in Madin-Darby canine kidney cells requires co-expressed human β ₂-microglobulin. *J. Biol. Chem.* 277: 28038-28050.
2. Praetor, A., et al. 2002. Membrane-anchored human FcRn can oligomerize in the absence of IgG. *J. Mol. Biol.* 321: 277-284.
3. Detmer, S.A., et al. 2002. IgG transcytosis and recycling by FcRn expressed in MDCK cells reveals ligand-induced redistribution. *EMBO J.* 21: 5953.
4. Zhou, J., et al. 2003. Generation of mutated variants of the human form of the MHC class I-related receptor, FcRn, with increased affinity for mouse immunoglobulin G. *J. Mol. Biol.* 332: 901-913.
6. Ober, R.J., et al. 2004. Exocytosis of IgG as mediated by the receptor, FcRn: an analysis at the single-molecule level. *Proc. Natl. Acad. Sci. USA* 101: 11076-11081.

CHROMOSOMAL LOCATION

Genetic locus: Fcgrt (mouse) mapping to 7 B4.

PRODUCT

FcRn siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FcRn shRNA Plasmid (m): sc-45633-SH and FcRn shRNA (m) Lentiviral Particles: sc-45633-V as alternate gene silencing products.

For independent verification of FcRn (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45633A, sc-45633B and sc-45633C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FcRn siRNA (m) is recommended for the inhibition of FcRn expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

FcRn (A-6): sc-393064 is recommended as a control antibody for monitoring of FcRn gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FcRn gene expression knockdown using RT-PCR Primer: FcRn (m)-PR: sc-45633-PR (20 μ l, 549 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Gan, H., et al. 2012. Neonatal Fc receptor stimulation induces ubiquitin C-terminal hydrolase-1 overexpression in podocytes through activation of p38 mitogen-activated protein kinase. *Hum. Pathol.* 43: 1482-1490.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.