



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mucin 13 shRNA (m) Lentiviral Particles: sc-45691-V

BACKGROUND

Mucins are epithelial glycoproteins with a high content of clustered oligosaccharides that are O-glycoside linked to tandem repeat peptides rich in threonine, serine and proline. Mucin 13 (MUC13), also designated down-regulated in colon cancer 1 (DRCC1), is an epithelial and hemopoietic type I membrane protein that undergoes secretion and influences gastrointestinal mucosa levels. It is most abundant in epithelial tissues of the gastrointestinal and respiratory tracts, such as large intestine and trachea, followed by kidney, small intestine, appendix and stomach. Mucin 13 is a good differentiation marker for gastrointestinal mucosa and may also indicate certain gastric tumors. It localizes to the apical membrane of both columnar and goblet cells in the gastrointestinal tract, and within goblet cell thecae. Mucin 13 is a cleaved protein, and the β subunit, containing the cytoplasmic tail, can homodimerize.

REFERENCES

- Williams, S.J., et al. 2001. Muc13, a novel human cell surface mucin expressed by epithelial and hemopoietic cells. *J. Biol. Chem.* 276: 18327-18336.
- Corrales, R.M., et al. 2003. Normal human conjunctival epithelium expresses MUC13, MUC15, MUC16 and MUC17 mucin genes. *Arch. Soc. Esp. Oftalmol.* 78: 375-381.
- Carraway, K.L., et al. 2003. Cell signaling through membrane mucins. *Bioessays* 25: 66-71.
- Packer, L.M., et al. 2004. Expression of the cell surface mucin gene family in adenocarcinomas. *Int. J. Oncol.* 25: 1119-1126.
- Byrd, J.C., et al. 2004. Mucins and mucin binding proteins in colorectal cancer. *Cancer Metastasis Rev.* 23: 77-99.
- Hollingsworth, M.A., et al. 2004. Mucins in cancer: protection and control of the cell surface. *Nat. Rev. Cancer* 4: 45-60.

CHROMOSOMAL LOCATION

Genetic locus: Muc13 (mouse) mapping to 16 B3.

PRODUCT

Mucin 13 shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Mucin 13 siRNA (m): sc-45691 and Mucin 13 shRNA Plasmid (m): sc-45691-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

Mucin 13 shRNA (m) Lentiviral Particles is recommended for the inhibition of Mucin 13 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Mucin 13 (G-10): sc-390115 is recommended as a control antibody for monitoring of Mucin 13 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Mucin 13 gene expression knockdown using RT-PCR Primer: Mucin 13 (m)-PR: sc-45691-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ$ C and the extension temperature should be $68-72^\circ$ C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.