

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Soggy-1 shRNA (m) Lentiviral Particles: sc-45756-V



The Power to Question

BACKGROUND

The Wnt genes, a group of well-conserved, cysteine-rich, secreted glycoproteins, are required for numerous developmental processes, including embryogenesis, asymmetric cell division and central nervous system (CNS) patterning. The association of the Wnt protein with the seven membrane spanning receptor frizzled activates dishevelled, which downregulates glycogen synthase kinase (GSK) through serine phosphorylation. Reduced levels of active GSK results in the accumulation of β -catenin and subsequent regulation of developmentally significant Wnt target genes. Wnt antagonists, which include Dickkopf (Dkk1-4), Soggy-1 (also designated DkkL1), frizzled-related protein (sFRP) and Wnt inhibitory factor-1 (WIF-1), are necessary to ensure normal spatial and temporal patterns of Wnt activity during developmental processes. Soggy-1 is a secreted antagonist of Wnt expressed in preimplantation mouse embryos and developing neural tissue, as well as adult testis tissue. Soggy-1 is subjected to posttranslational modification by glycosylation; additional modifications have been observed and are yet to be characterized. The molecular weight of Soggy-1 has been observed to vary depending on tissue and cell type.

REFERENCES

- Krasnow, R.E., et al. 1995. Dishevelled is a component of the frizzled signaling pathway in *Drosophila*. Development 121: 4095-4102.
- 2. Cadigan, K.M., et al. 1997. Wnt signaling: a common theme in animal development. Genes Dev. 11: 3286-3305.
- Sakanaka, C., et al. 1998. Bridging of β-catenin and glycogen synthase kinase-3β by axin and inhibition of β-catenin-mediated transcription. Proc. Natl. Acad. Sci. USA 95: 3020-3023.
- Glinka, A., et al. 1998. Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction. Nature 391: 357-362.
- 5. Krupnik, V.E., et al. 1999. Functional and structural diversity of the human Dickkopf gene family. Gene 238: 301-313.

CHROMOSOMAL LOCATION

Genetic locus: Dkkl1 (mouse) mapping to 7 B4.

PRODUCT

Soggy-1 shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0 x 10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Soggy-1 siRNA (m): sc-45756 and Soggy-1 shRNA Plasmid (m): sc-45756-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

Soggy-1 shRNA (m) Lentiviral Particles is recommended for the inhibition of Soggy-1 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0 x 10⁶ infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Soggy-1 (B-4): sc-393692 is recommended as a control antibody for monitoring of Soggy-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Soggy-1 gene expression knockdown using RT-PCR Primer: Soggy-1 (m)-PR: sc-45756-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**