



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DNAM-1 siRNA (m): sc-45776

BACKGROUND

The T cell antigen receptor (TCR) recognizes foreign antigens and translates such recognition events into intracellular signals that elicit a change in the cell from a dormant to an activated state. Additional proteins termed "accessory molecules" are also required for activation and for cytotoxic T lymphocyte (CTL)-mediated cytotoxicity. For instance, CD2, CD4, CD8, LFA-1, and CD28 are examples of well characterized accessory molecules. An accessory molecule designated DNAX accessory molecule-1 or DNAM-1 has been described. DNAM-1 is a transmembrane glycoprotein that is 318 amino acids in length and contains two immunoglobulin-like domains. DNAM-1 is expressed on both T cells and natural killer (NK) cells and participates in primary adhesion during CTL-mediated cytotoxicity.

REFERENCES

1. Arthos, J., et al. 1989. Identification of the residues in human CD4 critical for the binding of HIV. *Cell* 57: 469-481.
2. Weiss, A., et al. 1991. Signal transduction by the T cell antigen receptor. *Semin. Immunol.* 3: 313-324.
3. Allison, J.P., et al. 1991. The immunobiology of T cells with invariant γ δ antigen receptors. *Annu. Rev. Immunol.* 9: 679-705.
4. Ehrlich, E.W., et al. 1993. T cell receptor interaction with peptide/major histocompatibility complex (MHC) and superantigen/MHC ligands is dominated by antigen. *J. Exp. Med.* 178: 713-722.
5. Julius, M., et al. 1993. Distinct roles for CD4 and CD8 as co-receptors in antigen receptor signalling. *Immunol. Today* 14: 177-183.
6. Vignali, D.A. 1994. The interaction between CD4 and MHC class II molecules and its effect on T cell function. *Behring Inst. Mitt.* 94: 133-147.
7. Shibuya, A., et al. 1996. DNAM-1, A novel adhesion molecule involved in the cytolytic function of T lymphocytes. *Immunity* 4: 573-581.

CHROMOSOMAL LOCATION

Genetic locus: Cd226 (mouse) mapping to 18 E4.

PRODUCT

DNAM-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DNAM-1 shRNA Plasmid (m): sc-45776-SH and DNAM-1 shRNA (m) Lentiviral Particles: sc-45776-V as alternate gene silencing products.

For independent verification of DNAM-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45776A, sc-45776B and sc-45776C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DNAM-1 shRNA Plasmid (m) is recommended for the inhibition of DNAM-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DNAM-1 gene expression knockdown using RT-PCR Primer: DNAM-1 (m)-PR: sc-45776-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.