



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CHL1 siRNA (m): sc-45788

BACKGROUND

Neural cell adhesion molecules play an important role in neural cell interactions and neuronal development and growth. Close homolog of L1 (CHL1), also designated cell adhesion molecule with homology to L1CAM or L1 cell adhesion molecule 2, belongs to the L1 gene family of neural cell adhesion molecules. CHL1 plays a role in signal transduction pathways and is involved in cell migration, axon growth and guidance. It is required for neuronal positioning and dendritic growth of pyramidal neurons in the posterior region of the developing mouse neocortex. CHL1 is expressed in pyramidal neurons in a high-caudal to low-rostral gradient within the developing cortex.

REFERENCES

1. Angeloni, D., et al. 1999. CALL gene is haploinsufficient in a 3p-syndrome patient. *Am. J. Med. Genet.* 86: 482-485.
2. Sakurai, K., et al. 2002. An association between a missense polymorphism in the close homologue of L1 (CHL1, CALL) gene and schizophrenia. *Mol. Psychiatry* 7: 412-415.
3. Irintchev, A., et al. 2004. Impairment of sensorimotor gating in mice deficient in the cell adhesion molecule L1 or its close homologue, CHL1. *Brain Res.* 1029: 131-134.
4. Demyanenko, G.P., et al. 2004. CHL1 modulates area-specific neuronal positioning and dendrite orientation in the cerebral cortex. *Neuron* 44: 423-437.
5. Munos, S., et al. 2004. Transcript profiling in the CHL1-5 mutant of *Arabidopsis* reveals a role of the nitrate transporter NRT1.1 in the regulation of another nitrate transporter, NRT2.1. *Plant Cell* 16: 2433-2447.
6. Petronczki, M., et al. 2004. Sister-chromatid cohesion mediated by the alternative RF-CCTf18/Dcc1/Ctf8, the helicase CHL1 and the polymerase-alpha-associated protein Ctf4 is essential for chromatid disjunction during meiosis II. *J. Cell. Sci.* 117: 3547-3559.
7. Rokman, A., et al. 2005. Hereditary prostate cancer in Finland: fine-mapping validates 3p26 as a major predisposition locus. *Hum. Genet.* 116: 43-50.
8. Gast, D., et al. 2005. L1 augments cell migration and tumor growth but not Integrin β 3 expression in ovarian carcinomas. *Int. J. Cancer* 115: 658-665.

CHROMOSOMAL LOCATION

Genetic locus: Chl1 (mouse) mapping to 6 E1.

PRODUCT

CHL1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CHL1 shRNA Plasmid (m): sc-45788-SH and CHL1 shRNA (m) Lentiviral Particles: sc-45788-V as alternate gene silencing products.

For independent verification of CHL1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45788A, sc-45788B and sc-45788C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CHL1 siRNA (m) is recommended for the inhibition of CHL1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CHL1 gene expression knockdown using RT-PCR Primer: CHL1 (m)-PR: sc-45788-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.