



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HoxB7 siRNA (m): sc-45836

BACKGROUND

The Hox proteins play a role in development and cellular differentiation by regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. The mammalian HOX gene complex consists of 39 genes that are located on 4 linkage groups, which are dispersed over 4 chromosomes. A segment of the HoxB7 proximal promoter drives renal expression of reporter genes specifically in the ureteric bud and collecting ducts. Expression levels of HoxB7 are lower in lymph node metastasis-positive cancer tissues than negative cancer tissues. These results suggest that aberrant expression of HOX genes is related to the development of breast cancer and malignant behavior of cancer cells.

REFERENCES

1. Felicetti, F., et al. 2004. Role of PLZF in melanoma progression. *Oncogene* 23: 4567-4576.
2. Oxburgh, L., et al. 2004. TGF β superfamily signals are required for morphogenesis of the kidney mesenchyme progenitor population. *Development* 131: 4593-4605.
3. Watanabe, T. and Costantini, F. 2004. Real-time analysis of ureteric bud branching morphogenesis *in vitro*. *Dev. Biol.* 271: 98-108.
4. Yu, O.H., et al. 2004. Overexpression of Ret leads to vesicoureteric reflux in mice. *Am. J. Physiol. Renal Physiol.* 287: F1123-F1130.
5. Makiyama, K., et al. 2005. Aberrant expression of HOX genes in human invasive breast carcinoma. *Oncol. Rep.* 13: 673-679.
6. SWISS-PROT/TrEMBL (P09629). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: Hoxb7 (mouse) mapping to 11 D.

PRODUCT

HoxB7 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HoxB7 shRNA Plasmid (m): sc-45836-SH and HoxB7 shRNA (m) Lentiviral Particles: sc-45836-V as alternate gene silencing products.

For independent verification of HoxB7 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45836A, sc-45836B and sc-45836C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HoxB7 shRNA (m) Lentiviral Particles is recommended for the inhibition of HoxB7 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HoxB7 gene expression knockdown using RT-PCR Primer: HoxB7 (m)-PR: sc-45836-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.