



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



myomesin-1 shRNA (m) Lentiviral Particles: sc-45890-V

BACKGROUND

Myomesin-1 and myomesin-2 are components of the vertebrate myofibrillar M band and are associated with Titin, Myosin and Connectin. The myomesin proteins are responsible for the formation of a head structure on one end of the Titin string that connects the Z and M bands of the sarcomere. Myomesin-1 and -2 have unique N-terminal domains and are expressed mainly in skeletal muscle. The gene encoding human myomesin-1 maps to chromosome 18p11.31-p11.32.

REFERENCES

- Grove, B.K., et al. 1984. A new 185,000-dalton skeletal muscle protein detected by monoclonal antibodies. *J. Cell Biol.* 98: 518-524.
- Vinkemeier, U., et al. 1993. The globular head domain of titin extends into the center of the sarcomeric M band. cDNA cloning, epitope mapping and immunoelectron microscopy of two Titin-associated proteins. *J. Cell Sci.* 106: 319-330.
- Speel, E.J., et al. 1998. Assignment of the human gene for the sarcomeric M-band protein myomesin (MYOM1) to 18p11.31-p11.32. *Genomics* 54: 184-186.
- Agarkova, I., et al. 2000. A novel marker for vertebrate embryonic heart, the EH-myomesin isoform. *J. Biol. Chem.* 275: 10256-10264.
- Porter, J.D., et al. 2003. Postnatal suppression of myomesin, muscle creatine kinase and the M-line in rat extraocular muscle. *J. Exp. Biol.* 206: 3101-3112.
- Hornemann, T., et al. 2003. Muscle-type creatine kinase interacts with central domains of the M-band proteins myomesin and M-protein. *J. Mol. Biol.* 332: 877-887.

CHROMOSOMAL LOCATION

Genetic locus: Myom1 (mouse) mapping to 17 E1.3.

PRODUCT

myomesin-1 shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see myomesin-1 siRNA (m): sc-45890 and myomesin-1 shRNA Plasmid (m): sc-45890-SH as alternate gene silencing products.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

APPLICATIONS

myomesin-1 shRNA (m) Lentiviral Particles is recommended for the inhibition of myomesin-1 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

myomesin-1 (N-14): sc-30388 is recommended as a control antibody for monitoring of myomesin-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor myomesin-1 gene expression knockdown using RT-PCR Primer: myomesin-1 (m)-PR: sc-45890-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.