



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HIF-1 α siRNA (r): sc-45919

BACKGROUND

Cell growth and viability is compromised by oxygen deprivation (hypoxia). Hypoxia-inducible factors, including HIF-1 α , HIF-1 β (also designated Arnt 1), EPAS-1 (also designated HIF-2 α) and HIF-3 α , induce glycolysis, erythropoiesis and angiogenesis in order to restore oxygen homeostasis. Hypoxia-inducible factors are members of the Per-Arnt-Sim (PAS) domain transcription factor family. In response to hypoxia, HIF-1 α is upregulated and forms a heterodimer with Arnt 1 to form the HIF-1 complex. The HIF-1 complex recognizes and binds to the hypoxia responsive element (HRE) of hypoxia-inducible genes, thereby activating transcription. Hypoxia-inducible expression of some genes, such as Glut-1, p53, p21 or Bcl-2, is HIF-1 α dependent, whereas expression of others, such as p27, GADD 153 or HO-1, is HIF-1 α independent. EPAS-1 and HIF-3 α have also been shown to form heterodimeric complexes with Arnt 1 in response to hypoxia.

REFERENCES

1. Wang, G.L., et al. 1995. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc. Natl. Acad. Sci. USA* 92: 5510-5514.
2. Tian, H., et al. 1997. Endothelial PAS domain protein 1 (EPAS1), a transcription factor selectively expressed in endothelial cells. *Genes Dev.* 11: 72-82.
3. Luo, G., et al. 1997. Molecular characterization of the murine HIF-1 α locus. *Gene Expr.* 6: 287-299.

CHROMOSOMAL LOCATION

Genetic locus: Hif1a (rat) mapping to 6q24.

PRODUCT

HIF-1 α siRNA (r) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HIF-1 α shRNA Plasmid (r): sc-45919-SH and HIF-1 α shRNA (r) Lentiviral Particles: sc-45919-V as alternate gene silencing products.

For independent verification of HIF-1 α (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45919A, sc-45919B and sc-45919C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HIF-1 α siRNA (r) is recommended for the inhibition of HIF-1 α expression in rat cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

HIF-1 α (28b): sc-13515 is recommended as a control antibody for monitoring of HIF-1 α gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HIF-1 α gene expression knockdown using RT-PCR Primer: HIF-1 α (r)-PR: sc-45919-PR (20 μ l, 427 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

SELECT PRODUCT CITATIONS

1. Zhang, S., et al. 2007. Evidence of oxidative stress-induced BNIP3 expression in β -Amyloid neurotoxicity. *Brain Res.* 1138: 221-230.
2. Ryou M.G., et al. 2012. Pyruvate protects the brain against ischemia-reperfusion injury by activating the erythropoietin signaling pathway. *Stroke* 43: 1101-1107.
3. Wang, Y., et al. 2013. Hypoxia-inducible factor-1 α and MAPK co-regulate activation of hepatic stellate cells upon hypoxia stimulation. *PLoS ONE* 8: e74051.
4. Wang, K., et al. 2013. Mipu1, a novel direct target gene, is involved in hypoxia inducible factor 1-mediated cytoprotection. *PLoS ONE* 8: e82827.
5. Herrera-Martínez, M., et al. 2013. Actin, RhoA, and Rab11 participation during encystment in *Entamoeba invadens*. *Biomed Res. Int.* 2013: 919345.
6. Shan, F., et al. 2014. HIF-1 α -induced up-regulation of miR-9 contributes to phenotypic modulation in pulmonary artery smooth muscle cells during hypoxia. *J. Cell. Physiol.* 229: 1511-1520.
7. Jin, X., et al. 2015. Brain protection against ischemic stroke using choline as a new molecular bypass treatment. *Acta Pharmacol. Sin.* 36: 1416-1425.
8. Shao, Y., et al. 2016. Mir-217 promotes inflammation and fibrosis in high glucose cultured rat glomerular mesangial cells via Sirt1/HIF-1 α signaling pathway. *Diabetes Metab. Res. Rev.* 32: 534-543.

RESEARCH USE

For research use only, not for use in diagnostic procedures.