



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CRP3 shRNA (m) Lentiviral Particles: sc-45933-V

BACKGROUND

Cysteine and glycine-rich protein 3 (CRP3), also known as cysteine-rich protein 3, CLP (cardiac LIM protein), MLP (muscle LIM protein), LMO4 or CMD1M, is an essential nuclear regulator of myogenic differentiation. CRP3 contains two LIM zinc-binding domains linked to short glycine-rich repeats and a potential nuclear localization signal. CRP3 is present in differentiated heart during early development and in a subset of other striated muscles during later stages. Defects in the gene encoding CRP3 (CSRP3) can cause dilated cardiomyopathy 1M (CMD1M), a disease characterized by reduced systolic function and cardiac dilation. Human CSRP3 maps to the gene locus 11p15.1.

REFERENCES

1. Fung, Y., et al. 1995. Mapping of a human LIM protein (CLP) to human chromosome 11p15.1 by fluorescence *in situ* hybridization. *Genomics* 28: 602-603.
2. Weiskirchen, R., et al. 1995. The cysteine-rich protein family of highly related LIM domain proteins. *J. Biol. Chem.* 270: 28946-28954.
3. Knöll, R., et al. 2002. The cardiac mechanical stretch sensor machinery involves a Z-disc complex that is defective in a subset of human dilated cardiomyopathy. *Cell* 111: 943-955.
4. Duan, L., et al. 2003. Expression of muscle LIM protein during early development in *Xenopus laevis*. *Int. J. Dev. Biol.* 4: 299-302.
5. Geier, C., et al. 2003. Mutations in the human muscle LIM protein gene in families with hypertrophic cardiomyopathy. *Circulation* 10: 1390-1395.
6. Lu, P.Y., et al. 2004. Muscle LIM protein promotes expression of the acetylcholine receptor γ -subunit gene cooperatively with the myogenin-E12 complex. *Cell. Mol. Life Sci.* 61: 2386-2392.
7. Heineke, J., et al. 2005. Attenuation of cardiac remodeling after myocardial infarction by muscle LIM protein-calcineurin signaling at the sarcomeric Z-disc. *Proc. Natl. Acad. Sci. USA* 102: 1655-1660.

CHROMOSOMAL LOCATION

Genetic locus: *Csrp3* (mouse) mapping to 7 B4.

PRODUCT

CRP3 shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see CRP3 siRNA (m): sc-45933 and CRP3 shRNA Plasmid (m): sc-45933-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80°C . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4°C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

CRP3 shRNA (m) Lentiviral Particles is recommended for the inhibition of CRP3 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

CRP3 (B-4): sc-393599 is recommended as a control antibody for monitoring of CRP3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CRP3 gene expression knockdown using RT-PCR Primer: CRP3 (m)-PR: sc-45933-PR (20 μ l, 612 bp). Annealing temperature for the primers should be $55-60^\circ\text{C}$ and the extension temperature should be $68-72^\circ\text{C}$.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.