



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# RNase L siRNA (m): sc-45966

## BACKGROUND

RNase L encodes a component of the interferon-regulated 2-5A system that functions in the antiviral and antiproliferative roles of interferons. Mutations in this gene have been associated with predisposition to prostate cancer and this gene is a candidate for the hereditary prostate cancer 1 (HPC-1) allele. Interferon treatment enhances levels of both RNase L and a group of synthetases that produce 5'-triphosphorylated, 2',5'-oligoadenylates (2-5A) from ATP. The role of the 2-5A system in the control of viral and cellular growth suggests that defects in the 2-5A-dependent RNase gene could result in reduced immunity to virus infections and cancer.

## REFERENCES

1. Wu, H., et al. 1998. Molecular cloning and expression of cDNA for human RNase H. *Antisense Nucleic Acid Drug Dev.* 8: 53-61.
2. Cerritelli, S., et al. 1998. Cloning, expression, and mapping of ribonucleases H of human and mouse related to bacterial RNase HI. *Genomics* 53: 300-307.
3. Demette, E., et al. 2002. Ribonuclease L proteolysis in peripheral blood mononuclear cells of chronic fatigue syndrome patients. *J. Biol. Chem.* 277: 35746-35751.
4. ten Asbroek, A., et al. 2002. Ribonuclease H1 maps to chromosome 2 and has at least three pseudogene loci in the human genome. *Genomics* 79: 818-823.
5. Nakazato, H., et al. 2003. Role of genetic polymorphisms of the RNase L gene on familial prostate cancer risk in a Japanese population. *Br. J. Cancer* 89: 691-696.
6. Silverman, R.H. 2003. Implications for RNase L in prostate cancer biology. *Biochemistry* 42: 1805-1812.
7. Li, G., et al. 2004. An apoptotic signaling pathway in the interferon antiviral response mediated by RNase L and c-Jun NH<sub>2</sub>-terminal kinase. *J. Biol. Chem.* 279: 1123-1131.
8. SWISS-PROT/TrEMBL (O60930). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

## CHROMOSOMAL LOCATION

Genetic locus: Rnasel (mouse) mapping to 1 G3.

## PRODUCT

RNase L siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RNase L shRNA Plasmid (m): sc-45966-SH and RNase L shRNA (m) Lentiviral Particles: sc-45966-V as alternate gene silencing products.

For independent verification of RNase L (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45966A, sc-45966B and sc-45966C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

RNase L shRNA (m) Lentiviral Particles is recommended for the inhibition of RNase L expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

RNase L (E-9): sc-74405 is recommended as a control antibody for monitoring of RNase L gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RNase L gene expression knockdown using RT-PCR Primer: RNase L (m)-PR: sc-45966-PR (20  $\mu$ l, 586 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.