

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

PI-9 siRNA (m): sc-152245



BACKGROUND

Serine proteinase inhibitors (serpins) function as regulators of serine proteinase activity in a variety of physiological processes. Proteinase inhibitor-9 (PI-9, also designated cytoplasmic antiproteinase 3, or CAP3) is a member of the Ovalbumin family of serpins that is expressed in placenta, lung and cytotoxic lymphocytes. PI-9 is a potent inhibitor of granzyme B and of granzyme B-mediated apoptosis, and is also an inhibitor of caspase-1 and, to a lesser extent, caspase-4 and caspase-8. Because granzyme B promotes DNA degradation and rapidly translocates to the nucleus to bind to a nuclear component, PI-9 is present in the nuclei of human cytotoxic cells, endothelial cells and epithelial cells. PI-9 is exported from nuclei via a leptomycin B-sensitive pathway, suggesting that the nucleocytoplasmic distribution of PI-9 involves a nonconventional nuclear import pathway and the export factor CRM1. Estrogen rapidly and strongly induces PI-9, which is an estrogen-regulated human gene. PI-9 expression is also upregulated in response to inflammatory stimuli. This upregulation protects cells from apoptosis induced by endogenously expressed or released granzyme B, particulary during target cell killling. In addition, PI-9 is expressed in a variety of human and murine tumors.

REFERENCES

- Dahlen, J.R., et al. 1997. Human proteinase inhibitor-9 (PI-9) is a potent inhibitor of subtilisin A. Biochem. Biophys. Res. Commun. 238: 329-333.
- Sun, J., et al. 1997. A new family of 10 murine Ovalbumin serpins includes two homologs of proteinase inhibitor-8 and two homologs of the granzyme B inhibitor (proteinase inhibitor-9). J. Biol. Chem. 272: 15434-15441.
- Sun, J., et al. 1998. A serpin gene cluster on human chromosone 6p25 contains PI-6, PI-9 and ELANH₂ which have a common structure almost identical to the 18q21 Ovalbumin serpin genes. Cytogenet. Cell Genet. 82: 273-277.
- 4. Dahlen, J.R., et al. 1999. Inhibition of neutrophil elastase by recombinant human proteinase inhibitor-9. Biochem. Biophys. Acta 1451: 233-241.
- 5. Annand, R.R., et al. 1999. Caspase-1 (Interleukin-1 β -converting enzyme) is inhibited by the human serpin analogue proteinase inhibitor-9. Biochem. J. 342: 655-665.

CHROMOSOMAL LOCATION

Genetic locus: Serpinb9 (mouse) mapping to 13 A3.3.

PRODUCT

PI-9 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PI-9 shRNA Plasmid (m): sc-152245-SH and PI-9 shRNA (m) Lentiviral Particles: sc-152245-V as alternate gene silencing products.

For independent verification of PI-9 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152245A, sc-152245B and sc-152245C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

 $\mathsf{PI-9}$ siRNA (m) is recommended for the inhibition of $\mathsf{PI-9}$ expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

PI-9 (F-6): sc-390501 is recommended as a control antibody for monitoring of PI-9 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PI-9 gene expression knockdown using RT-PCR Primer: PI-9 (m)-PR: sc-152245-PR (20 μ I). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.