



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PI31 siRNA (m): sc-152248

BACKGROUND

Proteasomes are the catalytic sites of intracellular protein degradation via proteolysis. The most common form of proteasome, the 26S Proteasome, is composed of a cylindrical-shaped 20S core particle accompanied by two 19S regulatory subunits. PI31 (proteasome inhibitor PI31 subunit) is a 271 amino acid protein that is a potent inhibitor of proteasome activity. PI31 is a proline-rich protein, particularly within its C-terminal domain, where nearly one-quarter of the amino acids are proline. Working within the ubiquitin-dependent pathway, PI31 inhibits the hydrolysis of protein and peptide substrates by the 20S proteasome and also inhibits proteasome activation by PA700 and PA28. PI31 functions as a monomer and localizes to the nuclear envelope and endoplasmic reticulum membrane.

REFERENCES

1. Zaiss, D.M., et al. 1999. The proteasome inhibitor PI31 competes with PA28 for binding to 20S proteasomes. *FEBS Lett.* 457: 333-338.
2. McCutchen-Maloney, S.L., et al. 2000. cDNA cloning, expression, and functional characterization of PI31, a proline-rich inhibitor of the proteasome. *J. Biol. Chem.* 275: 18557-18565.
3. Zaiss, D.M., et al. 2002. PI31 is a modulator of proteasome formation and antigen processing. *Proc. Natl. Acad. Sci. USA* 99: 14344-14349.
4. Dahlmann, B. 2005. Proteasomes. *Essays Biochem.* 41: 31-48.
5. Gharbi, S.I., et al. 2007. Exploring the specificity of the PI3K family inhibitor LY294002. *Biochem. J.* 404: 15-21.
6. Kuznetsova, E.B., et al. 2007. Novel methylation and expression markers associated with breast cancer. *Mol. Biol.* 41: 624-633.
7. Kirk, R., et al. 2008. Structure of a conserved dimerization domain within the F-box protein Fbxo7 and the PI31 proteasome inhibitor. *J. Biol. Chem.* 283: 22325-22335.
8. Cui, Y., et al. 2008. Proteomic analysis of testis biopsies in men treated with injectable testosterone undecanoate alone or in combination with oral levonorgestrel as potential male contraceptive. *J. Proteome Res.* 7: 3984-3993.

CHROMOSOMAL LOCATION

Genetic locus: Psmf1 (mouse) mapping to 2 G3.

PRODUCT

PI31 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PI31 shRNA Plasmid (m): sc-152248-SH and PI31 shRNA (m) Lentiviral Particles: sc-152248-V as alternate gene silencing products.

For independent verification of PI31 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152248A, sc-152248B and sc-152248C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PI31 siRNA (m) is recommended for the inhibition of PI31 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PI31 gene expression knockdown using RT-PCR Primer: PI31 (m)-PR: sc-152248-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.