



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PPWD1 siRNA (m): sc-152430

BACKGROUND

PPWD1 (peptidylprolyl isomerase domain and WD repeat containing 1), also known as spliceosome-associated cyclophilin, is a 646 amino acid protein that belongs to the cyclophilin-type PPIase family and the PPIL1 subfamily. PPWD1 is similar in structure to canonical CypA, as well as to CypH and PPIL1, two additional spliceosomal cyclophilins. PPWD1 contains one PPIase cyclophilin-type domain and four WD repeats. Localizing to nucleus, PPWD1 associates with spliceosomal complexes and operates as a subunit of the spliceosome C complex. Functioning as a putative peptidylprolyl isomerase, PPWD1 accelerates protein folding and catalyzes the *cis-trans* isomerization of proline imidic peptide bonds in oligopeptides. PPWD1 may also play a role in pre-mRNA splicing. Highly conserved, PPWD1 is encoded by a gene that maps to human chromosome 5q12.3.

REFERENCES

1. Davis, T.L., Walker, J.R., Ouyang, H., MacKenzie, F., Butler-Cole, C., Newman, E.M., Eisenmesser, E.Z. and Dhe-Paganon, S. 2008. The crystal structure of human WD40 repeat-containing peptidylprolyl isomerase (PPWD1). *FEBS J.* 275: 2283-2295.
2. Mesa, A., Somarelli, J.A. and Herrera, R.J. 2008. Spliceosomal immunophilins. *FEBS Lett.* 582: 2345-2351.
3. Thai, V., Renesto, P., Fowler, C.A., Brown, D.J., Davis, T., Gu, W., Pollock, D.D., Kern, D., Raoult, D. and Eisenmesser, E.Z. 2008. Structural, biochemical, and *in vivo* characterization of the first virally encoded cyclophilin from the Mimivirus. *J. Mol. Biol.* 37: 71-86.
4. Bessonov, S., Anokhina, M., Will, C.L., Urlaub, H. and Lührmann, R. 2008. Isolation of an active step I spliceosome and composition of its RNP core. *Nature* 452: 846-850.
5. Kanlaya, R., Pattanakitsakul, S.N., Sinchaikul, S., Chen, S.T. and Thongboonkerd, V. 2010. Vimentin interacts with heterogeneous nuclear ribonucleoproteins and dengue nonstructural protein 1 and is important for viral replication and release. *Mol. Biosyst.* 6: 795-806.
6. Davis, T.L., Walker, J.R., Campagna-Slater, V., Finerty, P.J., Paramanathan, R., Bernstein, G., MacKenzie, F., Tempel, W., Ouyang, H., Lee, W.H., Eisenmesser, E.Z. and Dhe-Paganon, S. 2010. Structural and biochemical characterization of the human cyclophilin family of peptidyl-prolyl isomerases. *PLoS Biol.* 8: e1000439.
7. Bessonov, S., Anokhina, M., Krasauskas, A., Golas, M.M., Sander, B., Will, C.L., Urlaub, H., Stark, H. and Lührmann, R. 2010. Characterization of purified human Bact spliceosomal complexes reveals compositional and morphological changes during spliceosome activation and first step catalysis. *RNA* 16: 2384-2403.

CHROMOSOMAL LOCATION

Genetic locus: Ppwd1 (mouse) mapping to 13 D1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

PPWD1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PPWD1 shRNA Plasmid (m): sc-152430-SH and PPWD1 shRNA (m) Lentiviral Particles: sc-152430-V as alternate gene silencing products.

For independent verification of PPWD1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152430A, sc-152430B and sc-152430C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PPWD1 siRNA (m) is recommended for the inhibition of PPWD1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PPWD1 gene expression knockdown using RT-PCR Primer: PPWD1 (m)-PR: sc-152430-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.