



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

# PRAP1 siRNA (m): sc-152439

## BACKGROUND

PRAP1 (proline-rich acidic protein 1), also known as UPA (uterine-specific proline-rich acidic protein), is a 151 amino acid secreted protein that contains two putative casein kinase II phosphorylation sites and exists as four alternatively spliced isoforms. Human PRAP1 shares approximately 50% homology with both rat and mouse orthologues and contains a 20-amino acid signal peptide that is more than 70% conserved between human and rodent. Abundantly expressed in epithelial cells of liver, kidney, gastrointestinal tract and cervix, PRAP1 may play an important role in maintaining normal growth homeostasis in epithelial cells. PRAP1 is significantly down-regulated in hepatocellular carcinoma and right colon adenocarcinoma, compared to adjacent normal tissues, and conversely up-regulated by butyrate, Trichostatin A and 5'-aza-2' deoxycytidine. The gene that encodes PRAP1 maps to human chromosome 10q26.3.

## REFERENCES

1. Kasik, J. and Rice, E. 1997. A novel complementary deoxyribonucleic acid is abundantly and specifically expressed in the uterus during pregnancy. *Am. J. Obstet. Gynecol.* 176: 452-456.
2. Zhang, J., Rajkumar, N. and Hooi, S.C. 2000. Characterization and expression of the mouse pregnant specific uterus protein gene and its rat homologue in the intestine and uterus. *Biochim. Biophys. Acta* 1492: 526-530.
3. Zhang, J., Wong, H., Ramanan, S., Cheong, D., Leong, A. and Hooi, S.C. 2003. The proline-rich acidic protein is epigenetically regulated and inhibits growth of cancer cell lines. *Cancer Res.* 63: 6658-6665.
4. Laurin, N., Misener, V.L., Crosbie, J., Ickowicz, A., Pathare, T., Roberts, W., Malone, M., Tannock, R., Schachar, R., Kennedy, J.L. and Barr, C.L. 2005. Association of the calcyon gene (DRD1IP) with attention deficit/hyperactivity disorder. *Mol. Psychiatry* 10: 1117-1125.
5. Lanemo Myhrinder, A., Hellqvist, E., Sidorova, E., Söderberg, A., Baxendale, H., Dahle, C., Willander, K., Tobin, G., Bäckman, E., Söderberg, O., Rosenquist, R., Hörkö, S. and Rosen, A. 2008. A new perspective: molecular motifs on oxidized LDL, apoptotic cells, and bacteria are targets for chronic lymphocytic leukemia antibodies. *Blood* 111: 3838-3848.
6. Saito, S., Ojima, H., Ichikawa, H., Hirohashi, S. and Kondo, T. 2008. Molecular background of  $\alpha$ -fetoprotein in liver cancer cells as revealed by global RNA expression analysis. *Cancer Sci.* 99: 2402-2409.
7. Diao, H., Xiao, S., Zhao, F. and Ye, X. 2010. Uterine luminal epithelium-specific proline-rich acidic protein 1 (PRAP1) as a marker for successful embryo implantation. *Fertil. Steril.* 94: 2808-2811.e1.
8. Xiong, G.F., Zhang, Y.S., Han, B.C., Chen, W., Yang, Y. and Peng, J.P. 2011. Estradiol-regulated proline-rich acid protein 1 is repressed by class I histone deacetylase and functions in peri-implantation mouse uterus. *Mol. Cell. Endocrinol.* 331: 23-33.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Prap1 (mouse) mapping to 7 F4.

## PRODUCT

PRAP1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PRAP1 shRNA Plasmid (m): sc-152439-SH and PRAP1 shRNA (m) Lentiviral Particles: sc-152439-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

PRAP1 siRNA (m) is recommended for the inhibition of PRAP1 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PRAP1 gene expression knockdown using RT-PCR Primer: PRAP1 (m)-PR: sc-152439-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.