



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CENP-V siRNA (m): sc-152513

BACKGROUND

During mitosis, the transient assembly of the kinetochore occurs on a platform known as the centromere, a specialized chromatin structure that is comprised of various centromere proteins (CENPs). There are two multi-protein centromere complexes, known as CENPA-NAC (nucleosome-associated) and CENPA-CAD (nucleosome distal), which interact with one another to facilitate both the assembly and the activity of the centromere. CENP-V (centromere protein V), also known as PRR6 or nuclear protein p30, is a 275 amino acid protein that belongs to the gfa family. Existing as three alternatively spliced isoforms, CENP-V plays an essential role in cytokinesis, chromosome alignment, and the distribution of pericentromeric heterochromatin in interphase nuclei. The gene encoding CENP-V maps to human chromosome 17p11.2 and mouse chromosome 11 B2.

REFERENCES

1. Cronshaw, J.M., Krutchinsky, A.N., Zhang, W., Chait, B.T. and Matunis, M.J. 2002. Proteomic analysis of the mammalian nuclear pore complex. *J. Cell Biol.* 158: 915-927.
2. Izuta, H., Ikeno, M., Suzuki, N., Tomonaga, T., Nozaki, N., Obuse, C., Kisu, Y., Goshima, N., Nomura, F., Nomura, N. and Yoda, K. 2006. Comprehensive analysis of the ICEN (Interphase Centromere Complex) components enriched in the CENP-A chromatin of human cells. *Genes Cells* 11: 673-684.
3. Okada, M., Cheeseman, I.M., Hori, T., Okawa, K., McLeod, I.X., Yates, J.R., Desai, A. and Fukagawa, T. 2006. The CENP-H-I complex is required for the efficient incorporation of newly synthesized CENP-A into centromeres. *Nat. Cell Biol.* 8: 446-457.
4. Foltz, D.R., Jansen, L.E., Black, B.E., Bailey, A.O., Yates, J.R. and Cleveland, D.W. 2006. The human CENP-A centromeric nucleosome-associated complex. *Nat. Cell Biol.* 8: 458-469.
5. Tadeu, A.M., Ribeiro, S., Johnston, J., Goldberg, I., Gerloff, D. and Earnshaw, W.C. 2008. CENP-V is required for centromere organization, chromosome alignment and cytokinesis. *EMBO J.* 27: 2510-2522.
6. Honda, Z., Suzuki, T. and Honda, H. 2009. Identification of CENP-V as a novel microtubule-associating molecule that activates Src family kinases through SH3 domain interaction. *Genes Cells* 14: 1383-1394.

CHROMOSOMAL LOCATION

Genetic locus: Cenpv (mouse) mapping to 11 B2.

PRODUCT

CENP-V siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CENP-V shRNA Plasmid (m): sc-152513-SH and CENP-V shRNA (m) Lentiviral Particles: sc-152513-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CENP-V siRNA (m) is recommended for the inhibition of CENP-V expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CENP-V gene expression knockdown using RT-PCR Primer: CENP-V (m)-PR: sc-152513-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.