



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PSMD12 siRNA (m): sc-152558

BACKGROUND

In eukaryotic cells, selective breakdown of cellular proteins is ensured by their ubiquitination and subsequent degradation by the 26S Proteasome. The 26S Proteasome is a protease complex that selectively breaks down proteins that have been modified by polyubiquitin chains. It is made up of two multi-subunit complexes: the 20S Proteasome chamber, which serves as the proteolytic core of the complex and two 19S regulatory particles which recognize and unfold ubiquitinated proteins. PSMD12 (proteasome (prosome, macropain) 26S subunit, non-ATPase, 12), also known as p55 or Rpn5, is a 456 amino acid protein belonging to the proteasome subunit p55 family. PSMD12 acts as a regulatory subunit of the 26S Proteasome and is a component of the PA700 complex. PSMD12 contains one PCI domain.

REFERENCES

1. Yokota, K., et al. 1996. CDNA cloning of p112, the largest regulatory subunit of the human 26S Proteasome, and functional analysis of its yeast homologue, sen3p. *Mol. Biol. Cell* 7: 853-870.
2. Zhang, N., et al. 2003. Using yeast to place human genes in functional categories. *Gene* 303: 121-129.
3. Shibahara, T., et al. 2004. Mass spectrometric analysis of expression of ATPase subunits encoded by duplicated genes in the 19S regulatory particle of rice 26S Proteasome. *Arch. Biochem. Biophys.* 421: 34-41.
4. Thompson, H.G., et al. 2004. Post-translationally modified S12, absent in transformed breast epithelial cells, is not associated with the 26S Proteasome and is induced by proteasome inhibitor. *Int. J. Cancer* 111: 338-347.
5. Tan, Y., et al. 2006. Effects of tumor necrosis factor- α on the 26S Proteasome and 19S regulator in skeletal muscle of severely scalded mice. *J. Burn Care Res.* 27: 226-233.
6. Wang, Z., et al. 2006. Prostaglandin J2 alters pro-survival and pro-death gene expression patterns and 26S Proteasome assembly in human neuroblastoma cells. *J. Biol. Chem.* 281: 21377-21386.
7. Cui, F., et al. 2006. The up-regulation of proteasome subunits and lysosomal proteases in hepatocellular carcinomas of the HBx gene knockin transgenic mice. *Proteomics* 6: 498-504.

CHROMOSOMAL LOCATION

Genetic locus: Psmid12 (mouse) mapping to 11 E1.

PRODUCT

PSMD12 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PSMD12 shRNA Plasmid (m): sc-152558-SH and PSMD12 shRNA (m) Lentiviral Particles: sc-152558-V as alternate gene silencing products.

For independent verification of PSMD12 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152558A, sc-152558B and sc-152558C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PSMD12 siRNA (m) is recommended for the inhibition of PSMD12 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

PSMD12 (H-3): sc-398279 is recommended as a control antibody for monitoring of PSMD12 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PSMD12 gene expression knockdown using RT-PCR Primer: PSMD12 (m)-PR: sc-152558-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.